

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Home & Community Garden Soil Sample Survey

NEW YORK CITY

*Bronx, Queens, Kings,
Richmond and Manhattan*



Samples analyzed by the
Cornell Nutrient Analysis Laboratory (CNAL)
in 1995-2001

Quirine M. Ketterings, Hettie Krol, Charlie Mazza and W. Shaw Reid

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Home & Community Garden Soil Sample Survey

NEW YORK CITY

*Bronx, Queens, Kings,
Richmond and Manhattan*

Samples analyzed by CNAL in 1995-2001

Summary compiled by

Quirine Ketterings and Hettie Krol

Nutrient Management Spear Program, Department of Crop and Soil Sciences
817 Bradfield Hall, Cornell University, Ithaca NY 14853

Charles P. Mazza

Senior Extension Associate
Department of Horticulture

W. Shaw Reid

Professor Emeritus
Department of Crop and Soil Sciences

Ithaca, May 12, 2004

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Table of Content

| | |
|---|----|
| 1. Introduction to New York City Gardens and Soils..... | 1 |
| 2. General Survey Summary..... | 3 |
| 3. Crops/Plants | 13 |
| 3.1 Bronx | 13 |
| 3.2 Queens | 14 |
| 3.3 Kings | 15 |
| 3.4 Richmond | 16 |
| 3.5 Manhattan | 17 |
| 4. Soil Types | 18 |
| 4.1 Bronx | 18 |
| 4.2 Queens | 18 |
| 4.3 Kings | 19 |
| 4.4 Richmond | 19 |
| 4.5 Manhattan | 19 |
| 5. Organic Matter | 20 |
| 5.1 Bronx | 20 |
| 5.2 Queens | 21 |
| 5.3 Kings | 22 |
| 5.4 Richmond | 23 |
| 5.5 Manhattan | 24 |
| 6. pH | 25 |
| 6.1 Bronx | 25 |
| 6.2 Queens | 26 |
| 6.3 Kings | 27 |
| 6.4 Richmond | 28 |
| 6.5 Manhattan | 29 |
| 7. Phosphorus..... | 30 |
| 7.1 Bronx | 30 |
| 7.2 Queens | 31 |
| 7.3 Kings | 32 |

| | |
|------------------------------------|----|
| 7.4 Richmond | 33 |
| 7.5 Manhattan | 34 |
| 8. Potassium | 35 |
| 8.1 Bronx | 35 |
| 8.2 Queens | 38 |
| 8.3 Kings | 41 |
| 8.4 Richmond | 44 |
| 8.5 Manhattan | 47 |
| 9. Magnesium | 50 |
| 9.1 Bronx | 50 |
| 9.2 Queens | 51 |
| 9.3 Kings | 52 |
| 9.4 Richmond | 53 |
| 9.5 Manhattan | 54 |
| 10. Iron | 55 |
| 10.1 Bronx | 55 |
| 10.2 Queens | 56 |
| 10.3 Kings | 57 |
| 10.4 Richmond | 58 |
| 10.5 Manhattan | 59 |
| 11. Manganese | 60 |
| 11.1 Bronx | 60 |
| 11.2 Queens | 61 |
| 11.3 Kings | 62 |
| 11.4 Richmond | 63 |
| 11.5 Manhattan | 64 |
| 12. Zinc | 65 |
| 12.1 Bronx | 65 |
| 12.2 Queens | 66 |
| 12.3 Kings | 67 |
| 12.4 Richmond | 68 |
| 12.5 Manhattan | 69 |
| Appendix: Cornell Crop Codes | 71 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

1. Introduction to New York City Gardens and Soils

New York City is made up of five political boroughs that are also counties. New York City is an archipelago of islands along the Atlantic Ocean, with only a small part (the Bronx) on the mainland, north of the islands. Manhattan and Richmond (Staten Island) are separate islands, while Kings (Brooklyn) and Queens are boroughs/counties on the larger Long Island. A ridge along the northern end of Staten Island, Brooklyn, Queens (and extends eastward on Long Island) marks the terminal moraine of the last glacial period, making the land south of this ridge an outwash plain. Prior to the rapid residential and commercial development of Brooklyn, Queens and Staten Island, large parts of the outwash plains formed good, flat agricultural land.

The native soils vary from sandy along the Atlantic Ocean coast to a wider range of soil types in the constructed areas. Much of Manhattan Island, for instance, consists of fill from other sites. Some of the fill extended the shoreline of the islands; in many cases, the fill came from excavating for subways and large buildings with deep foundations and lower level construction. Bedrock in Manhattan is deep enough to carry the weight of the skyscrapers in Lower Manhattan (near the Wall Street Financial District) and mid-town (near the Rockefeller Center and Times Square Area). The area in between (Greenwich Village and Chelsea) has much shallower bedrock, which disallows for the construction of tall skyscrapers in that area.

There is one educational farm in Queens (on the Nassau County border), and a working farm as part of the State Parks System on Staten Island (Richmond Co.). In recent years, there has been some large-scale farming on Riker's Island (prison) – a portion of Queens. However, most edible and non-edible plants are grown in home gardens and community gardens. Home gardens abound in all boroughs, but especially in residential neighborhoods in Kings, Queens, Bronx and Richmond Counties. Community gardens – publicly owned land used to grow plants by groups of people – exist in all five boroughs (counties) of New York City but tend to be most abundant in areas where land is less desirable for construction and land values are depressed. The areas where many of the community gardens were established were often rubble strewn (brick, mortar, asphalt pieces, etc.) from remnants of former buildings. In many cases, where community gardens

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

are permitted for group use, new soil is brought in from outside New York City and poured into constructed raised beds. Issues concerning heavy metals from lead paint and other debris on these formerly abandoned sites have prompted caution. Community gardens often, but not always, grow edible plants, which might be contaminated if lead or other heavy metals were absorbed by the plants

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

2. General Survey Summary

This survey summarizes the soil test results from community garden and home garden soil samples from the Bronx, Queens, Kings, Richmond and Manhattan counties submitted for analyses to the Cornell Nutrient Analysis Laboratory (CNAL) during 1995-2001. The total number of samples analyzed in these years in New York City and vicinity amounted to 141 (Bronx), 351 (Queens), 399 (Kings), 100 (Richmond) and 574 (Manhattan) resulting in a total of 1565 samples over the 7 year period (Table 1).

Table 1: Total number of home and community garden samples submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001.

| | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|-------|----------|-----------|-----------|----------|-----------|------------|
| 1995 | 44 | 31 | 59 | 2 | 56 | 192 (12) |
| 1996 | 12 | 45 | 63 | 35 | 38 | 193 (12) |
| 1997 | 37 | 56 | 74 | 7 | 29 | 203 (13) |
| 1998 | 27 | 31 | 49 | 6 | 91 | 204 (13) |
| 1999 | 2 | 55 | 51 | 19 | 86 | 213 (14) |
| 2000 | 12 | 106 | 61 | 1 | 193 | 373 (24) |
| 2001 | 7 | 27 | 42 | 30 | 81 | 187 (12) |
| Total | 141 (9%) | 351 (22%) | 399 (26%) | 100 (6%) | 574 (37%) | 1565 (100) |

Twenty-two percent of the samples identified ornamentals adapted to slightly acid to calcareous soil conditions (SAG) as the target plants. Fifteen percent were submitted to obtain soil fertility data and recommendations for lawns while 10% of the samples came from home vegetable gardens (Table 2). A third of the samples that were submitted were sent in for lawns, perennials and mixed vegetable gardens. Eleven percent of the samples were submitted for perennials, 9% for ornamentals specially adapted to low pH (acidic) soils, and 4% came from athletic fields. The remainder of the samples was accompanied by requests for recommendations for parks, flowering annuals, roses, fruit trees, and other plants including berries, grapes, herbs, tomatoes, roadsides, etc. Summarizing the data, 12% of all samples were submitted to obtain recommendations for edible plants while

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

71% of the samples were submitted for recommendations for growing non-edible plants. For 17% of the samples, the plant to be grown was unknown to the laboratory.

Table 2: Number of samples submitted for soil fertility analyses and recommendations per crop/plant grown in home or community gardens in NYC in 1995-2001.

| | | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|-------|--------------------------|-------|--------|-------|----------|-----------|------------|
| ALG | Ornamentals (pH 4.5-6.0) | 6 | 28 | 26 | 0 | 74 | 134 (9) |
| ATF | Athletic Field | 1 | 6 | 15 | 28 | 5 | 55 (4) |
| BLU | Blueberries | 1 | 3 | 1 | 3 | 0 | 8 (<1) |
| FLA | Flowering Annuals | 8 | 8 | 15 | 7 | 19 | 57 (4) |
| GRA | Grapes | 0 | 1 | 1 | 0 | 0 | 2 (<1) |
| HRB | Herbs | 2 | 0 | 2 | 0 | 1 | 5 (<1) |
| IDL | Idle land | 0 | 2 | 0 | 0 | 5 | 7 (<1) |
| LAW | Lawn | 18 | 69 | 57 | 32 | 51 | 227 (15) |
| MVG | Mixed vegetables | 15 | 50 | 53 | 16 | 28 | 162 (10) |
| OTH | Other | 5 | 99 | 18 | 1 | 75 | 198 (13) |
| PER | Perennials | 13 | 25 | 66 | 9 | 60 | 173 (11) |
| PRK | Park | 2 | 1 | 5 | 0 | 75 | 83 (5) |
| ROD | Roadside | 0 | 2 | 3 | 0 | 1 | 6 (<1) |
| ROS | Roses | 5 | 2 | 6 | 0 | 17 | 30 (2) |
| RSP | Raspberries | 0 | 1 | 0 | 0 | 0 | 1 (<1) |
| SAG | Ornamentals(pH 6.0-7.5) | 49 | 34 | 118 | 3 | 136 | 340 (22) |
| SPB | Spring flowering bulbs | 4 | 0 | 3 | 0 | 2 | 9 (<1) |
| STR | Strawberries | 1 | 0 | 0 | 0 | 0 | 1 (<1) |
| SUB | Summer flowering bulbs | 0 | 0 | 0 | 0 | 1 | 1 (<1) |
| TOM | Tomatoes | 0 | 0 | 1 | 0 | 0 | 1 (<1) |
| TRF | Tree fruits | 1 | 0 | 2 | 0 | 4 | 7 (<1) |
| ? | Unknown (not specified) | 10 | 20 | 7 | 1 | 20 | 58 (4) |
| Total | | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Home and community garden samples from New York City were mostly silt loams (36%) and sandy loam soils (31%) belonging to soil management group 3 and 4, respectively (see Table 3 for a more detailed description of the soil management groups and Table 4 for a distribution of samples among the soil management groups). Eleven percent belonged to soil management group 2 which contains the silty soils. Group 5 (sandy soils)

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

was represented by 22% of all samples. There were no samples that were classified as clay or muck soils.

Table 3: Soil management groups for New York State.

| | |
|---|--|
| 1 | Fine-textured soils developed from clayey lake sediments and medium- to fine-textured soils developed from lake sediments. |
| 2 | Medium- to fine-textured soils developed from calcareous glacial till and medium-textured to moderately fine-textured soils developed from slightly calcareous glacial till mixed with shale and medium-textured soils developed in recent alluvium. |
| 3 | Moderately coarse textured soil developed from glacial outwash and recent alluvium and medium-textured acid soil developed on glacial till. |
| 4 | Coarse- to medium-textured soils formed from glacial till or glacial outwash. |
| 5 | Coarse- to very coarse-textured soils formed from gravelly or sandy glacial outwash or glacial lake beach ridges or deltas. |
| 6 | Organic or muck soils with more than 80% organic matter. |

Table 4: Number of home and community garden samples per soil management group (SMG) submitted to the Cornell Nutrient Analysis Laboratory during 1995-2001. The soil management groups (Table 3) impact availability of potassium and the recommendations.

| SMG | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|-------|-------|--------|-------|----------|-----------|------------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 (0) |
| 2 | 12 | 26 | 56 | 17 | 64 | 175 (11) |
| 3 | 40 | 140 | 149 | 59 | 176 | 564 (36) |
| 4 | 69 | 68 | 139 | 17 | 187 | 480 (31) |
| 5 | 20 | 117 | 55 | 7 | 147 | 346 (22) |
| 6 | 0 | 0 | 0 | 0 | 0 | 0 (0) |
| Total | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Organic matter levels of the samples varied widely. Twenty-three percent of the samples had between 3 and 4% organic matter while 13% had organic matter levels between 2 and 3% and 15% tested between 4 and 5% organic matter (Table 5). Organic matter levels greater than 5% were found in 32% of the samples while 17% contained less than 2% organic matter.

Table 5: Distribution of samples per organic matter level for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory during 1995-2001.

| Organic Matter (%) | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|--------------------|-------|--------|-------|----------|-----------|------------|
| <1 | 2 | 78 | 10 | 1 | 54 | 145 (9) |
| 1.0-1.9 | 7 | 36 | 35 | 5 | 42 | 125 (8) |
| 2.0-2.9 | 19 | 47 | 68 | 16 | 53 | 203 (13) |
| 3.0-3.9 | 35 | 68 | 87 | 32 | 131 | 353 (23) |
| 4.0-4.9 | 32 | 43 | 69 | 14 | 82 | 240 (15) |
| 5.0-5.9 | 15 | 27 | 46 | 4 | 61 | 153 (10) |
| 6.0-6.9 | 8 | 18 | 27 | 1 | 36 | 90 (6) |
| >6.9 | 23 | 34 | 57 | 27 | 115 | 256 (16) |
| Total | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Soil pH is a measure of soil acidity. Some plants are adapted to lower pH while others grow best on higher pH soils (generally pH 6 and over). Table 6 shows examples of ornamentals adapted to low versus higher pH status.

The pH values of the soils submitted to the Cornell Nutrient Analysis Laboratory varied widely (Table 7). Six percent of the samples had pH less than 5.0 (Table 6). Twenty-four percent tested between pH 5 and pH 6 while pH values over 6 but less than 8 were found for 66% of the samples. Really high pH values of 8 and higher (calcareous soils) were found for 3% of the samples.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Table 6: Ornamentals adapted pH less than or greater than pH 6.0.

| | |
|-----------------------|---|
| Adapted to pH 4.5-6.0 | Azalea, Bayberry, Bigleaf Hydrangea, Chokeberry, Franklinia, Holly, Inkberry, Leucothoe, Mountain Laurel, Oak, Pachistima, Pieris, Rhododendron, Sheep Laurel, Sourwood, Spicebush, Winterberry |
| Adapted to pH 6.0-7.5 | Abelia, Almond, Ajuga, Arborvitae, Ash, Barberry, Beautybush, Birch (White), Bittersweet, Boxwood, Chastetree, Chestnut, Clematis, Coralberry, Cotoneaster, Crabapple, Cranberry bush, Cypress, Daphne, Deutiza, Dogwood, Enkianthus, Euonymus, Firethorn, Fir, Forsythia, Fringe Tree, Germanander, Ginko, Golden Chain, Hawthorn, Hemlock, Hollygrape, Honey Locust, Honeysuckle, Hornbeam, Hydrangea, Hypericum, Ivy, Jetbead, Juniper, Larch, Lilac, Linden, Magnolia, Maple, Mockorange, Oak (English, Scarlet, Turkey), Pea Shrub, Pine, Plum (Flowering), Privet, Quince, Redbud, Rose of Sharon, Sassafras, Spirea, Spruce, Sweet Gum, Sweet Shrub, Sycamore, Tulip Tree, Tupelo (Gum), Va. Creeper, Viburnum, Vinca, Walnut, Wayfaring Tree, Weigela, Willow, Wisteria, Witch Hazel, Yellow-wood, Yew. |

Table 7: Number of samples in each of the pH classes for home and community garden soils submitted between 1995 and 2001.

| pH | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|---------|-------|--------|-------|----------|-----------|------------|
| <4.5 | 2 | 4 | 1 | 1 | 9 | 17 (1) |
| 4.5-4.9 | 16 | 14 | 9 | 1 | 43 | 83 (5) |
| 5.0-5.4 | 15 | 34 | 24 | 7 | 59 | 139 (9) |
| 5.5-5.9 | 19 | 75 | 39 | 16 | 79 | 228 (15) |
| 6.0-6.4 | 23 | 60 | 75 | 21 | 94 | 273 (17) |
| 6.5-6.9 | 17 | 77 | 114 | 20 | 124 | 352 (23) |
| 7.0-7.4 | 32 | 51 | 83 | 22 | 109 | 297 (19) |
| 7.5-7.9 | 9 | 28 | 42 | 11 | 43 | 133 (9) |
| 8.0-8.4 | 2 | 6 | 10 | 1 | 11 | 30 (2) |
| >8.4 | 6 | 2 | 2 | 0 | 3 | 13 (<1) |
| Total | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Extractable nutrients such as phosphorus (P), potassium (K), magnesium (Mg), iron (Fe), manganese (Mn), and zinc (Zn) were measured using the Morgan chemical extraction solution and method. This solution contains sodium acetate buffered at a pH of 4.8. Other extraction methods exist that give very different results.

Soil test phosphorus levels of <1 lb P/acre are classified as very low. Between 1-3 lbs P/acre is low. Medium is between 4-8 lbs P/acre. High testing soils have P levels between 9 and 39 lbs P/acre and soils with 40 lbs P/acre or more are classified as very high.

Of the home and community garden samples from New York City that were submitted to the Cornell Nutrient Analysis Laboratory between 1995 and 2001, none tested very low in phosphorus (Table 8). Five percent of the samples tested low in phosphorus while 8% were classified medium and 32% tested high in P. Fifty-four percent of the samples tested very high in phosphorus. **This meant that for 86% of the soils that were tested, for most plants, no additional phosphorus fertilizer would be needed.**

Table 8: Distribution of samples over different soil phosphorus availability classes. Soil test phosphorus levels of <1 lb P/acre are classified as very low. Between 1-3 lbs P/acre is low. Medium is between 4-8 lbs P/acre. High testing soils have P levels between 9 and 39 lbs P/acre and soils with 40 lbs P/acre or more are classified as very high.

| Soil Test lbs P/acre | Classification | Bronx | Queens | Kings | Rich- mond | Man- hattan | Total (%) |
|-------------------------|----------------|-------|--------|-------|---------------|----------------|------------|
| <1 | Very Low | 0 | 0 | 0 | 0 | 0 | 0 (0) |
| 1-3 | Low | 9 | 31 | 4 | 7 | 22 | 73 (5) |
| 4-8 | Medium | 11 | 45 | 14 | 21 | 41 | 132 (8) |
| 9-39 | High | 59 | 145 | 115 | 25 | 151 | 495 (32) |
| 40-60 | Very High | 29 | 44 | 105 | 8 | 99 | 285 (18) |
| 61-80 | Very High | 9 | 21 | 41 | 3 | 55 | 129 (8) |
| 81-100 | Very High | 5 | 16 | 20 | 2 | 45 | 88 (6) |
| 101-150 | Very High | 5 | 25 | 36 | 6 | 61 | 133 (9) |
| 151-200 | Very High | 5 | 9 | 25 | 2 | 28 | 69 (4) |
| >200 | Very High | 9 | 15 | 39 | 26 | 72 | 161 (10) |
| Total | | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Classifications for potassium in soils throughout the entire state of New York depend on soil management group. The fine-textured soils of soil management group 1 contain a lot of potassium-containing clay and have as a result a greater K supplying capacity than the coarse-textured sandy soils of soil management group 5. Because of these differences in potassium supplying capacity among soils of different origins (soil management groups as outlined in Table 3), the classifications and interpretations for potassium availability differ among the six groups. This is shown in Table 9. So for example for soils in soil management group 3, <45 lbs K/acre in the soil test means the soil is very low in K. If the soil test is between 45 and 79 lbs K/acre the soil is classified as low in potassium. Between 80 and 119 lbs K/acre is considered medium; between 120 and 199 lbs K/acre is high, and >199 lbs K/acre is classified as very high in plant available potassium (Table 9).

Table 9: Potassium soil test interpretations for New York State soils.

| Soil Management Group | Cornell Potassium Soil Test (Morgan extraction in lbs K/acre) | | | | |
|-----------------------|---|--------|---------|---------|-----------|
| | Very Low | Low | Medium | High | Very High |
| 1 | <35 | 35-64 | 65-94 | 95-149 | >149 |
| 2 | <40 | 40-69 | 70-99 | 100-164 | >164 |
| 3 | <45 | 45-79 | 80-119 | 120-199 | >199 |
| 4 | <55 | 55-99 | 100-149 | 150-239 | >239 |
| 5 and 6 | <60 | 60-114 | 115-164 | 165-269 | >269 |

Of the home and community garden samples submitted from New York City during 1995-2001, 9% were classified as very low in potassium (Table 10). Six percent had low potassium availability while 11% were classified as medium in potassium. High potassium availability was identified in 22% of the samples whereas 52% of the samples were classified as very high in potassium. **Thus, 74% of the samples tested high or very high for potassium where for many crops and non-edible plants only small amounts or none additional potassium would be recommended.**

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Table 10: Number of samples in each of the potassium soil fertility classes for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001. The classification (low, medium, etc.) of the actual soil tests depend on the specific soil and its soil management group (see Tables 3 and 9).

| Classification | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|----------------|-------|--------|-------|----------|-----------|------------|
| Very Low | 1 | 66 | 9 | 1 | 59 | 136 (9) |
| Low | 4 | 50 | 20 | 6 | 19 | 99 (6) |
| Medium | 23 | 40 | 60 | 13 | 35 | 171 (11) |
| High | 43 | 70 | 91 | 24 | 124 | 352 (23) |
| Very High | 70 | 125 | 219 | 56 | 337 | 807 (52) |
| Total | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Soils test very low for magnesium if Morgan extractable Mg is less than 20 lbs Mg/acre. Low testing soils have 20-65 lbs Morgan Mg per acre. Soils with 66-100 lbs Mg/acre test medium for magnesium. High testing soils have 101-199 lbs Mg/acre while soils with more than 200 lbs Mg/acre in the Morgan extraction are classified as very high in Mg. A little over one percent of the samples tested very low in magnesium. **Most soils tested high (15%) or very high (72%) while 8% tested low and 3% tested medium. Thus, magnesium deficiency is unlikely to occur for most soils in New York City**

Table 11: Number of samples in each of the magnesium soil fertility classes for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001.

| Soil test lbs Mg/acre | Classification | Bronx | Queens | Kings | Rich- mond | Man- hattan | Total (%) |
|--------------------------|----------------|-------|--------|-------|---------------|----------------|------------|
| <20 | Very low | 2 | 15 | 0 | 0 | 3 | 20 (1) |
| 20-65 | Low | 6 | 47 | 10 | 1 | 57 | 121 (8) |
| 66-100 | Medium | 7 | 17 | 10 | 3 | 17 | 54 (4) |
| 101-199 | High | 15 | 60 | 72 | 6 | 83 | 236 (15) |
| 200+ | Very high | 111 | 212 | 307 | 90 | 414 | 1134 (72) |
| Total | | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Soils with more than 50 lbs Morgan extractable iron per acre test excessive for iron. Anything lower than 50 lbs Fe/acre is considered normal. Of the 1565 samples, 1467 (94%) were classified as normal in iron availability (Table 12). The remainder had more iron than needed for optimum plant growth and were hence classified as excessive in iron.

Table 12: Number of samples testing normal or excessive for iron for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001.

| Soil test lbs Fe/acre | Classification | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|--------------------------|----------------|-------|--------|-------|----------|-----------|------------|
| <50 | Normal | 102 | 332 | 379 | 96 | 558 | 1467 (94) |
| 50+ | Excessive | 39 | 19 | 20 | 4 | 16 | 98 (6) |
| Total | Total | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Soils with more than 100 lbs Morgan extractable manganese per acre are classified as excessive in Mn. Anything less than 100 lbs Mn per acre is classified as normal. Of the 1565 samples that were submitted, 1520 (97%) were classified as normal in manganese availability (Table 13). The remainder of the samples had more manganese than needed for optimum plant growth and were hence classified as excessive in manganese.

Table 13: Number of samples testing normal or excessive for manganese for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001.

| Soil test lbs Mn/acre | Classification | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|--------------------------|----------------|-------|--------|-------|----------|-----------|------------|
| <100 | Normal | 130 | 349 | 391 | 84 | 566 | 1520 (97) |
| 100+ | Excessive | 11 | 2 | 8 | 16 | 8 | 45 (3) |
| Total | | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Soils with less than 0.5 lb zinc per acre in the Morgan extraction are classified as low in Zn. Medium testing soils have between 0.5 and 1 lb of Morgan extractable Zn per acre. If more than 1 lb of Zn/acre is extracted with the Morgan solution, the soil tests high in Zn. **For the home and community garden samples, 98% tested high for zinc while less than 2% tested medium in zinc (Table 14).**

Table 14: Number of samples testing low, medium or high in zinc for iron for home and community garden samples from New York City submitted to the Cornell Nutrient Analysis Laboratory in 1995-2001.

| Soil test lbs Zn/acre | Classification | Bronx | Queens | Kings | Richmond | Manhattan | Total (%) |
|--------------------------|----------------|-------|--------|-------|----------|-----------|------------|
| <0.5 | Low | 0 | 0 | 0 | 0 | 0 | 0 (0) |
| 0.5-1.0 | Medium | 0 | 10 | 3 | 1 | 10 | 24 (2) |
| >1.0 | High | 141 | 341 | 396 | 99 | 564 | 1541 (98) |
| Total | | 141 | 351 | 399 | 100 | 574 | 1565 (100) |

In the following sections, the summary tables for each of the soil fertility indicators described above are given. The appendix contains the crop codes used in section 3.

3. Crops/Plants

3.1 Bronx

Plants/crops for which recommendations are requested by homeowners:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total | % |
|---------|------|------|------|------|------|------|------|-------|-----|
| ALG | 0 | 0 | 2 | 3 | 0 | 1 | 0 | 6 | 4 |
| ATF | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 |
| BLU | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 1 |
| FLA | 0 | 3 | 2 | 0 | 2 | 1 | 0 | 8 | 6 |
| HRB | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 1 |
| LAW | 2 | 1 | 12 | 1 | 0 | 2 | 0 | 18 | 13 |
| MVG | 1 | 5 | 3 | 1 | 0 | 1 | 4 | 15 | 11 |
| OTH | 0 | 0 | 1 | 4 | 0 | 0 | 0 | 5 | 4 |
| PER | 0 | 0 | 8 | 1 | 0 | 3 | 1 | 13 | 9 |
| PRK | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 2 | 1 |
| ROS | 4 | 0 | 1 | 0 | 0 | 0 | 0 | 5 | 4 |
| SAG | 37 | 1 | 3 | 5 | 0 | 3 | 0 | 49 | 35 |
| SPB | 0 | 0 | 3 | 0 | 0 | 0 | 1 | 4 | 3 |
| STR | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 1 |
| TRF | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 1 |
| Unknown | 0 | 0 | 0 | 10 | 0 | 0 | 0 | 10 | 7 |
| | | | | | | | | | |
| Total | 44 | 12 | 37 | 27 | 2 | 12 | 7 | 141 | 100 |

Notes:

See Appendix for Cornell crop codes.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

3.2 Queens

Crops for which recommendations are requested by homeowners:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total | % |
|---------|------|------|------|------|------|------|------|-------|-----|
| ALG | 12 | 2 | 8 | 0 | 1 | 5 | 0 | 28 | 8 |
| ATF | 0 | 0 | 0 | 1 | 1 | 2 | 2 | 6 | 2 |
| BLU | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 3 | 1 |
| FLA | 0 | 1 | 2 | 1 | 2 | 2 | 0 | 8 | 2 |
| GRA | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| IDL | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 2 | 1 |
| LAW | 5 | 19 | 8 | 13 | 7 | 13 | 4 | 69 | 20 |
| MVG | 6 | 17 | 4 | 5 | 2 | 15 | 1 | 50 | 14 |
| OTH | 3 | 0 | 25 | 0 | 13 | 55 | 3 | 99 | 28 |
| PER | 1 | 1 | 0 | 2 | 3 | 9 | 9 | 25 | 7 |
| PRK | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 |
| ROD | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 2 | 1 |
| ROS | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 2 | 1 |
| RSP | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| SAG | 2 | 4 | 4 | 7 | 11 | 3 | 3 | 34 | 10 |
| Unknown | 0 | 0 | 0 | 0 | 14 | 1 | 5 | 20 | 6 |
| Total | 31 | 45 | 56 | 31 | 55 | 106 | 27 | 351 | 100 |

Notes:

See Appendix for Cornell crop codes.

3.3 Kings

Crops for which recommendations are requested by homeowners:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total | % |
|---------|------|------|------|------|------|------|------|-------|-----|
| ALG | 2 | 3 | 3 | 9 | 6 | 3 | 0 | 26 | 7 |
| ATF | 0 | 0 | 0 | 2 | 10 | 2 | 1 | 15 | 4 |
| BLU | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 |
| FLA | 1 | 5 | 4 | 3 | 1 | 1 | 0 | 15 | 4 |
| GRA | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 |
| HRB | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | 1 |
| LAW | 14 | 3 | 13 | 6 | 6 | 4 | 11 | 57 | 14 |
| MVG | 7 | 13 | 13 | 5 | 3 | 4 | 8 | 53 | 13 |
| OTH | 1 | 3 | 1 | 1 | 3 | 9 | 0 | 18 | 5 |
| PER | 3 | 6 | 21 | 9 | 9 | 12 | 6 | 66 | 17 |
| PRK | 4 | 0 | 0 | 1 | 0 | 0 | 0 | 5 | 1 |
| ROD | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 3 | 1 |
| ROS | 0 | 2 | 3 | 1 | 0 | 0 | 0 | 6 | 2 |
| SAG | 25 | 26 | 11 | 11 | 11 | 22 | 12 | 118 | 30 |
| SPB | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 3 | 1 |
| TOM | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 |
| TRF | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 2 | 1 |
| Unknown | 2 | 0 | 1 | 0 | 0 | 4 | 0 | 7 | 2 |
| Total | 59 | 63 | 74 | 49 | 51 | 61 | 42 | 399 | 100 |

Notes:

See Appendix for Cornell crop codes.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

3.4 Richmond

Crops for which recommendations are requested by homeowners:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total | % |
|---------|------|------|------|------|------|------|------|-------|-----|
| ATF | 0 | 28 | 0 | 0 | 0 | 0 | 0 | 28 | 28 |
| BLU | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 3 | 3 |
| FLA | 0 | 0 | 0 | 0 | 2 | 0 | 5 | 7 | 7 |
| LAW | 1 | 2 | 1 | 4 | 5 | 0 | 19 | 32 | 32 |
| MVG | 1 | 3 | 4 | 1 | 2 | 1 | 4 | 16 | 16 |
| OTH | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 |
| PER | 0 | 1 | 1 | 1 | 6 | 0 | 0 | 9 | 9 |
| SAG | 0 | 1 | 0 | 0 | 0 | 0 | 2 | 3 | 3 |
| Unknown | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | 1 |
| Total | 2 | 35 | 7 | 6 | 19 | 1 | 30 | 100 | 100 |

Notes:

See Appendix for Cornell crop codes.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

3.5 Manhattan

Crops for which recommendations are requested by homeowners:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total | % |
|---------|------|------|------|------|------|------|------|-------|-----|
| ALG | 6 | 4 | 3 | 9 | 21 | 23 | 8 | 74 | 13 |
| ATF | 0 | 0 | 0 | 1 | 0 | 4 | 0 | 5 | 1 |
| FLA | 4 | 1 | 2 | 4 | 2 | 0 | 6 | 19 | 3 |
| HRB | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| IDL | 0 | 0 | 0 | 0 | 5 | 0 | 0 | 5 | 1 |
| LAW | 3 | 3 | 3 | 7 | 4 | 11 | 20 | 51 | 9 |
| MVG | 9 | 2 | 1 | 3 | 5 | 4 | 4 | 28 | 5 |
| OTH | 1 | 1 | 0 | 2 | 8 | 57 | 6 | 75 | 13 |
| PER | 7 | 4 | 1 | 10 | 14 | 7 | 17 | 60 | 10 |
| PRK | 3 | 0 | 0 | 2 | 1 | 64 | 5 | 75 | 13 |
| ROD | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| ROS | 1 | 0 | 0 | 7 | 9 | 0 | 0 | 17 | 3 |
| SAG | 18 | 20 | 18 | 38 | 14 | 13 | 15 | 136 | 24 |
| SPB | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 | 0 |
| SUB | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 |
| TRF | 0 | 1 | 0 | 2 | 1 | 0 | 0 | 4 | 1 |
| Unknown | 3 | 1 | 1 | 6 | 2 | 7 | 0 | 20 | 3 |
| Total | 56 | 38 | 29 | 91 | 86 | 193 | 81 | 574 | 100 |

Notes:

See Appendix for Cornell crop codes.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

4. Soil Types

4.1 Bronx

Soil types (soil management groups) for home and garden samples:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total |
|--------------------|------|------|------|------|------|------|------|-------|
| SMG 1 (clayey) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| SMG 2 (silty) | 0 | 2 | 8 | 1 | 0 | 0 | 1 | 12 |
| SMG 3 (silt loam) | 1 | 2 | 15 | 16 | 0 | 5 | 1 | 40 |
| SMG 4 (sandy loam) | 43 | 5 | 7 | 6 | 0 | 7 | 1 | 69 |
| SMG 5 (sandy) | 0 | 3 | 7 | 4 | 2 | 0 | 4 | 20 |
| SMG 6 (mucky) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 44 | 12 | 37 | 27 | 2 | 12 | 7 | 141 |

4.2 Queens

Soil types (soil management groups) for home and garden samples:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total |
|--------------------|------|------|------|------|------|------|------|-------|
| SMG 1 (clayey) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| SMG 2 (silty) | 5 | 3 | 5 | 4 | 0 | 7 | 2 | 26 |
| SMG 3 (silt loam) | 5 | 20 | 9 | 2 | 44 | 51 | 9 | 140 |
| SMG 4 (sandy loam) | 5 | 20 | 12 | 13 | 5 | 5 | 8 | 68 |
| SMG 5 (sandy) | 16 | 2 | 30 | 12 | 6 | 43 | 8 | 117 |
| SMG 6 (mucky) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 31 | 45 | 56 | 31 | 55 | 106 | 27 | 351 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

4.3 Kings

Soil types (soil management groups) for home and garden samples:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total |
|--------------------|------|------|------|------|------|------|------|-------|
| SMG 1 (clayey) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| SMG 2 (silty) | 2 | 18 | 19 | 7 | 5 | 3 | 2 | 56 |
| SMG 3 (silt loam) | 25 | 15 | 16 | 26 | 25 | 31 | 11 | 149 |
| SMG 4 (sandy loam) | 26 | 18 | 32 | 9 | 11 | 17 | 26 | 139 |
| SMG 5 (sandy) | 6 | 12 | 7 | 7 | 10 | 10 | 3 | 55 |
| SMG 6 (mucky) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 59 | 63 | 74 | 49 | 51 | 61 | 42 | 399 |

4.4 Richmond

Soil types (soil management groups) for home and garden samples:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total |
|--------------------|------|------|------|------|------|------|------|-------|
| SMG 1 (clayey) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| SMG 2 (silty) | 1 | 5 | 1 | 1 | 5 | 0 | 4 | 17 |
| SMG 3 (silt loam) | 1 | 27 | 2 | 0 | 6 | 0 | 23 | 59 |
| SMG 4 (sandy loam) | 0 | 1 | 2 | 5 | 6 | 0 | 3 | 17 |
| SMG 5 (sandy) | 0 | 2 | 2 | 0 | 2 | 1 | 0 | 7 |
| SMG 6 (mucky) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 2 | 35 | 7 | 6 | 19 | 1 | 30 | 100 |

4.5 Manhattan

Soil types (soil management groups) for home and garden samples:

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | Total |
|--------------------|------|------|------|------|------|------|------|-------|
| SMG 1 (clayey) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| SMG 2 (silty) | 4 | 5 | 1 | 15 | 18 | 12 | 9 | 64 |
| SMG 3 (silt loam) | 31 | 13 | 10 | 19 | 30 | 50 | 23 | 176 |
| SMG 4 (sandy loam) | 15 | 12 | 12 | 38 | 21 | 55 | 34 | 187 |
| SMG 5 (sandy) | 6 | 8 | 6 | 19 | 17 | 76 | 15 | 147 |
| SMG 6 (mucky) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 56 | 38 | 29 | 91 | 86 | 193 | 81 | 574 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

5. Organic Matter

5.1 Bronx

Number of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 1 | 5 | 8 | 11 | 6 | 2 | 11 | 44 |
| 1996 | 0 | 1 | 0 | 3 | 5 | 1 | 1 | 1 | 12 |
| 1997 | 1 | 1 | 4 | 13 | 10 | 4 | 1 | 3 | 37 |
| 1998 | 1 | 1 | 5 | 3 | 4 | 3 | 3 | 7 | 27 |
| 1999 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 2 |
| 2000 | 0 | 3 | 2 | 4 | 1 | 0 | 1 | 1 | 12 |
| 2001 | 0 | 0 | 2 | 3 | 1 | 1 | 0 | 0 | 7 |
| Total | 2 | 7 | 19 | 35 | 32 | 15 | 8 | 23 | 141 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1.9 | 1.5 | 0.9 | 0.9 | 2.4 | 1.1 | 2.3 | |
| Highest: | 10.4 | 9.7 | 12.5 | 25.0 | 3.0 | 11.8 | 5.9 | |
| Mean: | 5.3 | 4.6 | 4.1 | 7.3 | 2.7 | 4.0 | 3.6 | |
| Median: | 4.6 | 4.4 | 3.9 | 4.9 | 2.7 | 3.5 | 3.1 | |

Percent of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 2 | 11 | 18 | 25 | 14 | 5 | 25 | 100 |
| 1996 | 0 | 8 | 0 | 25 | 42 | 8 | 8 | 8 | 100 |
| 1997 | 3 | 3 | 11 | 35 | 27 | 11 | 3 | 8 | 100 |
| 1998 | 4 | 4 | 19 | 11 | 15 | 11 | 11 | 26 | 100 |
| 1999 | 0 | 0 | 50 | 50 | 0 | 0 | 0 | 0 | 100 |
| 2000 | 0 | 25 | 17 | 33 | 8 | 0 | 8 | 8 | 100 |
| 2001 | 0 | 0 | 29 | 43 | 14 | 14 | 0 | 0 | 100 |
| Total | 1 | 5 | 13 | 25 | 23 | 11 | 6 | 16 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

5.2 Queens

Number of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 2 | 8 | 6 | 5 | 3 | 1 | 6 | 31 |
| 1996 | 1 | 1 | 8 | 9 | 8 | 3 | 7 | 8 | 45 |
| 1997 | 22 | 5 | 5 | 9 | 9 | 3 | 1 | 2 | 56 |
| 1998 | 0 | 6 | 5 | 12 | 3 | 3 | 1 | 1 | 31 |
| 1999 | 20 | 4 | 5 | 12 | 4 | 4 | 2 | 4 | 55 |
| 2000 | 34 | 16 | 13 | 12 | 11 | 8 | 3 | 9 | 106 |
| 2001 | 1 | 2 | 3 | 8 | 3 | 3 | 3 | 4 | 27 |
| Total | 78 | 36 | 47 | 68 | 43 | 27 | 18 | 34 | 351 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1.1 | 0.7 | 0.1 | 1.4 | 0.1 | 0.1 | 0.9 | |
| Highest: | 14.0 | 13.3 | 12.7 | 8.8 | 11.8 | 9.9 | 12.1 | |
| Mean: | 4.8 | 5.1 | 2.5 | 3.4 | 2.8 | 2.8 | 4.7 | |
| Median: | 3.9 | 4.4 | 2.1 | 3.2 | 2.7 | 2.1 | 3.9 | |

Percent of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 6 | 26 | 19 | 16 | 10 | 3 | 19 | 100 |
| 1996 | 2 | 2 | 18 | 20 | 18 | 7 | 16 | 18 | 100 |
| 1997 | 39 | 9 | 9 | 16 | 16 | 5 | 2 | 4 | 100 |
| 1998 | 0 | 19 | 16 | 39 | 10 | 10 | 3 | 3 | 100 |
| 1999 | 36 | 7 | 9 | 22 | 7 | 7 | 4 | 7 | 100 |
| 2000 | 32 | 15 | 12 | 11 | 10 | 8 | 3 | 8 | 100 |
| 2001 | 4 | 7 | 11 | 30 | 11 | 11 | 11 | 15 | 100 |
| Total | 22 | 10 | 13 | 19 | 12 | 8 | 5 | 10 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

5.3 Kings

Number of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 5 | 13 | 14 | 13 | 5 | 4 | 5 | 59 |
| 1996 | 2 | 11 | 5 | 7 | 11 | 8 | 6 | 13 | 63 |
| 1997 | 1 | 2 | 14 | 22 | 15 | 8 | 5 | 7 | 74 |
| 1998 | 1 | 3 | 7 | 9 | 5 | 10 | 3 | 11 | 49 |
| 1999 | 1 | 6 | 11 | 11 | 12 | 4 | 3 | 3 | 51 |
| 2000 | 4 | 7 | 12 | 9 | 7 | 7 | 5 | 10 | 61 |
| 2001 | 1 | 1 | 6 | 15 | 6 | 4 | 1 | 8 | 42 |
| Total | 10 | 35 | 68 | 87 | 69 | 46 | 27 | 57 | 399 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1.1 | 0.7 | 0.2 | 0.7 | 0.8 | 0.5 | 0.7 | |
| Highest: | 10.9 | 22.1 | 14.8 | 19.4 | 19.6 | 36.8 | 18.1 | |
| Mean: | 4.1 | 5.3 | 4.4 | 5.6 | 4.0 | 5.0 | 5.4 | |
| Median: | 3.8 | 4.7 | 3.9 | 4.9 | 3.7 | 3.7 | 3.8 | |

Percent of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 8 | 22 | 24 | 22 | 8 | 7 | 8 | 100 |
| 1996 | 3 | 17 | 8 | 11 | 17 | 13 | 10 | 21 | 100 |
| 1997 | 1 | 3 | 19 | 30 | 20 | 11 | 7 | 9 | 100 |
| 1998 | 2 | 6 | 14 | 18 | 10 | 20 | 6 | 22 | 100 |
| 1999 | 2 | 12 | 22 | 22 | 24 | 8 | 6 | 6 | 100 |
| 2000 | 7 | 11 | 20 | 15 | 11 | 11 | 8 | 16 | 100 |
| 2001 | 2 | 2 | 14 | 36 | 14 | 10 | 2 | 19 | 100 |
| Total | 3 | 9 | 17 | 22 | 17 | 12 | 7 | 14 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

5.4 Richmond

Number of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 2 |
| 1996 | 0 | 1 | 7 | 23 | 2 | 1 | 0 | 1 | 35 |
| 1997 | 0 | 0 | 0 | 1 | 2 | 1 | 0 | 3 | 7 |
| 1998 | 0 | 1 | 2 | 1 | 1 | 0 | 0 | 1 | 6 |
| 1999 | 1 | 0 | 3 | 5 | 6 | 1 | 0 | 3 | 19 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 3 | 4 | 2 | 3 | 0 | 0 | 18 | 30 |
| Total | 1 | 5 | 16 | 32 | 14 | 4 | 1 | 27 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 5.7 | 1.6 | 3.8 | 1.9 | 0.8 | 10.0 | 1.3 | |
| Highest: | 6.8 | 10.0 | 14.8 | 7.3 | 19.8 | 10.0 | 14.2 | |
| Mean: | 6.3 | 3.4 | 6.9 | 3.6 | 4.9 | 10.0 | 8.5 | |
| Median: | 6.3 | 3.2 | 5.7 | 2.9 | 4.2 | 10.0 | 10.5 | |

Percent of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | . | . | . | . | . | . | . | . | . |
| 1996 | . | . | . | . | . | . | . | . | . |
| 1997 | . | . | . | . | . | . | . | . | . |
| 1998 | . | . | . | . | . | . | . | . | . |
| 1999 | . | . | . | . | . | . | . | . | . |
| 2000 | . | . | . | . | . | . | . | . | . |
| 2001 | . | . | . | . | . | . | . | . | . |
| Total | 1 | 5 | 16 | 32 | 14 | 4 | 1 | 27 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

5.5 Manhattan

Number of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 3 | 7 | 8 | 12 | 11 | 5 | 2 | 8 | 56 |
| 1996 | 0 | 2 | 1 | 12 | 6 | 4 | 4 | 9 | 38 |
| 1997 | 0 | 3 | 3 | 9 | 3 | 1 | 6 | 4 | 29 |
| 1998 | 3 | 4 | 7 | 19 | 10 | 11 | 2 | 35 | 91 |
| 1999 | 7 | 5 | 8 | 15 | 15 | 6 | 5 | 25 | 86 |
| 2000 | 40 | 16 | 21 | 42 | 22 | 21 | 7 | 24 | 193 |
| 2001 | 1 | 5 | 5 | 22 | 15 | 13 | 10 | 10 | 81 |
| Total | 54 | 42 | 53 | 131 | 82 | 61 | 36 | 115 | 574 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 0.4 | 1.6 | 1.4 | 0.8 | 0.1 | 0.1 | 0.9 | |
| Highest: | 40.2 | 19.5 | 13.6 | 39.7 | 37.3 | 38.1 | 27.1 | |
| Mean: | 4.9 | 6.5 | 4.8 | 8.6 | 7.8 | 4.0 | 5.2 | |
| Median: | 3.9 | 4.8 | 3.9 | 5.2 | 4.3 | 3.5 | 4.6 | |

Percent of home and garden samples within each % organic matter range:

| | <1% | 1.0-1.9 | 2.0-2.9 | 3.0-3.9 | 4.0-4.9 | 5.0-5.9 | 6.0-6.9 | >6.9 | Total |
|-------|-----|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 5 | 13 | 14 | 21 | 20 | 9 | 4 | 14 | 100 |
| 1996 | 0 | 5 | 3 | 32 | 16 | 11 | 11 | 24 | 100 |
| 1997 | 0 | 10 | 10 | 31 | 10 | 3 | 21 | 14 | 100 |
| 1998 | 3 | 4 | 8 | 21 | 11 | 12 | 2 | 38 | 100 |
| 1999 | 8 | 6 | 9 | 17 | 17 | 7 | 6 | 29 | 100 |
| 2000 | 21 | 8 | 11 | 22 | 11 | 11 | 4 | 12 | 100 |
| 2001 | 1 | 6 | 6 | 27 | 19 | 16 | 12 | 12 | 100 |
| Total | 9 | 7 | 9 | 23 | 14 | 11 | 6 | 20 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

6. pH

6.1 Bronx

Number of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 2 | 12 | 10 | 7 | 5 | 0 | 5 | 2 | 0 | 1 | 44 |
| 1996 | 0 | 0 | 0 | 1 | 1 | 2 | 4 | 4 | 0 | 0 | 12 |
| 1997 | 0 | 0 | 1 | 5 | 9 | 8 | 8 | 0 | 1 | 5 | 37 |
| 1998 | 0 | 4 | 3 | 6 | 7 | 2 | 5 | 0 | 0 | 0 | 27 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 2 |
| 2000 | 0 | 0 | 1 | 0 | 1 | 4 | 4 | 1 | 1 | 0 | 12 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 1 | 6 | 0 | 0 | 0 | 7 |
| Total | 2 | 16 | 15 | 19 | 23 | 17 | 32 | 9 | 2 | 6 | 141 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 4.3 | 5.7 | 5.3 | 4.5 | 7.5 | 5.3 | 6.8 | |
| Highest: | 9.6 | 7.9 | 9.5 | 7.4 | 7.7 | 8.0 | 7.3 | |
| Mean: | - | - | - | - | - | - | - | |
| Median: | 5.3 | 7.2 | 6.6 | 6.1 | 7.6 | 6.9 | 7.1 | |

Percent of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 5 | 27 | 23 | 16 | 11 | 0 | 11 | 5 | 0 | 2 | 100 |
| 1996 | 0 | 0 | 0 | 8 | 8 | 17 | 33 | 33 | 0 | 0 | 100 |
| 1997 | 0 | 0 | 3 | 14 | 24 | 22 | 22 | 0 | 3 | 14 | 100 |
| 1998 | 0 | 15 | 11 | 22 | 26 | 7 | 19 | 0 | 0 | 0 | 100 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 100 |
| 2000 | 0 | 0 | 8 | 0 | 8 | 33 | 33 | 8 | 8 | 0 | 100 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 14 | 86 | 0 | 0 | 0 | 100 |
| Total | 1 | 11 | 11 | 13 | 16 | 12 | 23 | 6 | 1 | 4 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

6.2 Queens

Number of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 1 | 0 | 0 | 6 | 4 | 7 | 9 | 4 | 0 | 0 | 31 |
| 1996 | 0 | 4 | 5 | 5 | 10 | 10 | 4 | 5 | 1 | 1 | 45 |
| 1997 | 1 | 5 | 4 | 11 | 8 | 15 | 8 | 3 | 0 | 1 | 56 |
| 1998 | 0 | 1 | 3 | 6 | 5 | 11 | 5 | 0 | 0 | 0 | 31 |
| 1999 | 2 | 2 | 6 | 12 | 10 | 7 | 9 | 6 | 1 | 0 | 55 |
| 2000 | 0 | 2 | 15 | 30 | 19 | 17 | 13 | 6 | 4 | 0 | 106 |
| 2001 | 0 | 0 | 1 | 5 | 4 | 10 | 3 | 4 | 0 | 0 | 27 |
| Total | 4 | 14 | 34 | 75 | 60 | 77 | 51 | 28 | 6 | 2 | 351 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 4.3 | 4.7 | 4.3 | 4.5 | 4.4 | 4.7 | 5.3 | |
| Highest: | 7.8 | 8.7 | 9.3 | 7.3 | 8.0 | 8.4 | 7.9 | |
| Mean: | - | - | - | - | - | - | - | |
| Median: | 6.9 | 6.4 | 6.3 | 6.5 | 6.3 | 6.2 | 6.6 | |

Percent of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 3 | 0 | 0 | 19 | 13 | 23 | 29 | 13 | 0 | 0 | 100 |
| 1996 | 0 | 9 | 11 | 11 | 22 | 22 | 9 | 11 | 2 | 2 | 100 |
| 1997 | 2 | 9 | 7 | 20 | 14 | 27 | 14 | 5 | 0 | 2 | 100 |
| 1998 | 0 | 3 | 10 | 19 | 16 | 35 | 16 | 0 | 0 | 0 | 100 |
| 1999 | 4 | 4 | 11 | 22 | 18 | 13 | 16 | 11 | 2 | 0 | 100 |
| 2000 | 0 | 2 | 14 | 28 | 18 | 16 | 12 | 6 | 4 | 0 | 100 |
| 2001 | 0 | 0 | 4 | 19 | 15 | 37 | 11 | 15 | 0 | 0 | 100 |
| Total | 1 | 4 | 10 | 21 | 17 | 22 | 15 | 8 | 2 | 1 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

6.3 Kings

Number of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 1 | 5 | 7 | 12 | 24 | 6 | 3 | 1 | 0 | 59 |
| 1996 | 0 | 0 | 1 | 5 | 7 | 21 | 16 | 13 | 0 | 0 | 63 |
| 1997 | 1 | 3 | 8 | 10 | 15 | 22 | 14 | 0 | 0 | 1 | 74 |
| 1998 | 0 | 1 | 4 | 5 | 12 | 10 | 11 | 2 | 3 | 1 | 49 |
| 1999 | 0 | 0 | 3 | 7 | 9 | 9 | 13 | 10 | 0 | 0 | 51 |
| 2000 | 0 | 2 | 2 | 2 | 13 | 10 | 16 | 10 | 6 | 0 | 61 |
| 2001 | 0 | 2 | 1 | 3 | 7 | 18 | 7 | 4 | 0 | 0 | 42 |
| Total | 1 | 9 | 24 | 39 | 75 | 114 | 83 | 42 | 10 | 2 | 399 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 4.6 | 5.2 | 3.7 | 4.8 | 5.1 | 4.6 | 4.9 | |
| Highest: | 8.2 | 7.9 | 8.7 | 8.8 | 7.9 | 8.3 | 7.6 | |
| Mean: | - | - | - | - | - | - | - | |
| Median: | 6.6 | 6.9 | 6.5 | 6.5 | 6.7 | 7.0 | 6.7 | |

Percent of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 2 | 8 | 12 | 20 | 41 | 10 | 5 | 2 | 0 | 100 |
| 1996 | 0 | 0 | 2 | 8 | 11 | 33 | 25 | 21 | 0 | 0 | 100 |
| 1997 | 1 | 4 | 11 | 14 | 20 | 30 | 19 | 0 | 0 | 1 | 100 |
| 1998 | 0 | 2 | 8 | 10 | 24 | 20 | 22 | 4 | 6 | 2 | 100 |
| 1999 | 0 | 0 | 6 | 14 | 18 | 18 | 25 | 20 | 0 | 0 | 100 |
| 2000 | 0 | 3 | 3 | 3 | 21 | 16 | 26 | 16 | 10 | 0 | 100 |
| 2001 | 0 | 5 | 2 | 7 | 17 | 43 | 17 | 10 | 0 | 0 | 100 |
| Total | 0 | 2 | 6 | 10 | 19 | 29 | 21 | 11 | 3 | 1 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

6.4 Richmond

Number of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 2 |
| 1996 | 0 | 0 | 0 | 10 | 10 | 7 | 8 | 0 | 0 | 0 | 35 |
| 1997 | 0 | 0 | 0 | 0 | 1 | 3 | 2 | 1 | 0 | 0 | 7 |
| 1998 | 0 | 0 | 1 | 1 | 1 | 0 | 3 | 0 | 0 | 0 | 6 |
| 1999 | 1 | 1 | 1 | 0 | 3 | 4 | 1 | 7 | 1 | 0 | 19 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 2001 | 0 | 0 | 5 | 5 | 6 | 5 | 6 | 3 | 0 | 0 | 30 |
| Total | 1 | 1 | 7 | 16 | 21 | 20 | 22 | 11 | 1 | 0 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 6.5 | 5.5 | 6.4 | 5.2 | 4.3 | 7.1 | 5.1 | |
| Highest: | 7.4 | 7.4 | 7.9 | 7.4 | 8.3 | 7.1 | 7.7 | |
| Mean: | - | - | - | - | - | - | - | |
| Median: | 6.9 | 6.3 | 6.8 | 6.7 | 6.8 | 7.1 | 6.4 | |

Percent of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | . | . | . | . | . | . | . | . | . | . | . |
| 1996 | . | . | . | . | . | . | . | . | . | . | . |
| 1997 | . | . | . | . | . | . | . | . | . | . | . |
| 1998 | . | . | . | . | . | . | . | . | . | . | . |
| 1999 | . | . | . | . | . | . | . | . | . | . | . |
| 2000 | . | . | . | . | . | . | . | . | . | . | . |
| 2001 | . | . | . | . | . | . | . | . | . | . | . |
| Total | 1 | 1 | 7 | 16 | 21 | 20 | 22 | 11 | 1 | 0 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

6.5 Manhattan

Number of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 2 | 5 | 1 | 5 | 8 | 11 | 18 | 5 | 0 | 1 | 56 |
| 1996 | 1 | 2 | 3 | 10 | 7 | 7 | 1 | 3 | 4 | 0 | 38 |
| 1997 | 1 | 1 | 0 | 8 | 4 | 9 | 5 | 1 | 0 | 0 | 29 |
| 1998 | 2 | 8 | 6 | 12 | 18 | 21 | 14 | 8 | 2 | 0 | 91 |
| 1999 | 0 | 2 | 13 | 14 | 13 | 21 | 8 | 8 | 5 | 2 | 86 |
| 2000 | 2 | 21 | 26 | 24 | 30 | 34 | 43 | 13 | 0 | 0 | 193 |
| 2001 | 1 | 4 | 10 | 6 | 14 | 21 | 20 | 5 | 0 | 0 | 81 |
| Total | 9 | 43 | 59 | 79 | 94 | 124 | 109 | 43 | 11 | 3 | 574 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 4.3 | 4.4 | 4.4 | 4.2 | 4.8 | 4.1 | 4.4 | |
| Highest: | 9.5 | 8.2 | 7.5 | 8.4 | 9.3 | 7.8 | 7.9 | |
| Mean: | - | - | - | - | - | - | - | |
| Median: | 6.8 | 6.2 | 6.5 | 6.4 | 6.5 | 6.4 | 6.6 | |

Percent of home and garden samples within each pH range:

| | <4.5 | 4.5-4.9 | 5.0-5.4 | 5.5-5.9 | 6.0-6.4 | 6.5-6.9 | 7.0-7.4 | 7.5-7.9 | 8.0-8.4 | >8.4 | Total |
|-------|------|---------|---------|---------|---------|---------|---------|---------|---------|------|-------|
| 1995 | 4 | 9 | 2 | 9 | 14 | 20 | 32 | 9 | 0 | 2 | 100 |
| 1996 | 3 | 5 | 8 | 26 | 18 | 18 | 3 | 8 | 11 | 0 | 100 |
| 1997 | 3 | 3 | 0 | 28 | 14 | 31 | 17 | 3 | 0 | 0 | 100 |
| 1998 | 2 | 9 | 7 | 13 | 20 | 23 | 15 | 9 | 2 | 0 | 100 |
| 1999 | 0 | 2 | 15 | 16 | 15 | 24 | 9 | 9 | 6 | 2 | 100 |
| 2000 | 1 | 11 | 13 | 12 | 16 | 18 | 22 | 7 | 0 | 0 | 100 |
| 2001 | 1 | 5 | 12 | 7 | 17 | 26 | 25 | 6 | 0 | 0 | 100 |
| Total | 2 | 7 | 10 | 14 | 16 | 22 | 19 | 7 | 2 | 1 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

7. Phosphorus

7.1 Bronx

Number of home and garden samples within each range Morgan extractable P range (lbs/acre Morgan P):

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 8 | 5 | 20 | 3 | 3 | 0 | 1 | 1 | 3 | 44 |
| 1996 | 0 | 0 | 0 | 5 | 3 | 1 | 0 | 0 | 1 | 2 | 12 |
| 1997 | 0 | 0 | 0 | 17 | 8 | 5 | 3 | 1 | 1 | 2 | 37 |
| 1998 | 0 | 1 | 4 | 10 | 8 | 0 | 1 | 1 | 1 | 1 | 27 |
| 1999 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 2 |
| 2000 | 0 | 0 | 2 | 5 | 1 | 0 | 1 | 2 | 1 | 0 | 12 |
| 2001 | 0 | 0 | 0 | 2 | 4 | 0 | 0 | 0 | 0 | 1 | 7 |
| Total | 0 | 9 | 11 | 59 | 29 | 9 | 5 | 5 | 5 | 9 | 141 |

VL = very low, L = low, M = medium, H = high, VH = very high.

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1 | 24 | 9 | 3 | 48 | 6 | 22 | |
| Highest: | 301 | 483 | 438 | 212 | 51 | 194 | 204 | |
| Mean: | 45 | 112 | 64 | 43 | 50 | 60 | 68 | |
| Median: | 22 | 41 | 44 | 25 | 50 | 28 | 52 | |

Percent of home and garden samples within each Morgan extractable phosphorus range:

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 18 | 11 | 45 | 7 | 7 | 0 | 2 | 2 | 7 | 100 |
| 1996 | 0 | 0 | 0 | 42 | 25 | 8 | 0 | 0 | 8 | 17 | 100 |
| 1997 | 0 | 0 | 0 | 46 | 22 | 14 | 8 | 3 | 3 | 5 | 100 |
| 1998 | 0 | 4 | 15 | 37 | 30 | 0 | 4 | 4 | 4 | 4 | 100 |
| 1999 | 0 | 0 | 0 | 0 | 100 | 0 | 0 | 0 | 0 | 0 | 100 |
| 2000 | 0 | 0 | 17 | 42 | 8 | 0 | 8 | 17 | 8 | 0 | 100 |
| 2001 | 0 | 0 | 0 | 29 | 57 | 0 | 0 | 0 | 0 | 14 | 100 |
| Total | 0 | 6 | 8 | 42 | 21 | 6 | 4 | 4 | 4 | 6 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

7.2 Queens

Number of home and garden samples within each range Morgan extractable P range (lbs/acre Morgan P):

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 0 | 3 | 13 | 4 | 3 | 2 | 2 | 1 | 3 | 31 |
| 1996 | 0 | 0 | 0 | 17 | 6 | 3 | 5 | 7 | 2 | 5 | 45 |
| 1997 | 0 | 15 | 2 | 24 | 6 | 3 | 0 | 3 | 3 | 0 | 56 |
| 1998 | 0 | 0 | 6 | 14 | 5 | 3 | 0 | 1 | 1 | 1 | 31 |
| 1999 | 0 | 3 | 12 | 28 | 3 | 2 | 2 | 5 | 0 | 0 | 55 |
| 2000 | 0 | 9 | 18 | 43 | 17 | 5 | 7 | 5 | 0 | 2 | 106 |
| 2001 | 0 | 4 | 4 | 6 | 3 | 2 | 0 | 2 | 2 | 4 | 27 |
| Total | 0 | 31 | 45 | 145 | 44 | 21 | 16 | 25 | 9 | 15 | 351 |

VL = very low, L = low, M = medium, H = high, VH = very high.

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 6 | 13 | 1 | 4 | 3 | 1 | 2 | |
| Highest: | 835 | 495 | 169 | 221 | 143 | 285 | 490 | |
| Mean: | 88 | 92 | 34 | 42 | 30 | 37 | 86 | |
| Median: | 39 | 56 | 16 | 26 | 15 | 22 | 28 | |

Percent of home and garden samples within each Morgan extractable phosphorus range:

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 0 | 10 | 42 | 13 | 10 | 6 | 6 | 3 | 10 | 100 |
| 1996 | 0 | 0 | 0 | 38 | 13 | 7 | 11 | 16 | 4 | 11 | 100 |
| 1997 | 0 | 27 | 4 | 43 | 11 | 5 | 0 | 5 | 5 | 0 | 100 |
| 1998 | 0 | 0 | 19 | 45 | 16 | 10 | 0 | 3 | 3 | 3 | 100 |
| 1999 | 0 | 5 | 22 | 51 | 5 | 4 | 4 | 9 | 0 | 0 | 100 |
| 2000 | 0 | 8 | 17 | 41 | 16 | 5 | 7 | 5 | 0 | 2 | 100 |
| 2001 | 0 | 15 | 15 | 22 | 11 | 7 | 0 | 7 | 7 | 15 | 100 |
| Total | 0 | 9 | 13 | 41 | 13 | 6 | 5 | 7 | 3 | 4 | 100 |

VL = very low, L = low, M = medium, H = high, VH = very high.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

7.3 Kings

Number of home and garden samples within each range Morgan extractable P range (lbs/acre Morgan P):

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 1 | 0 | 21 | 14 | 9 | 2 | 10 | 1 | 1 | 59 |
| 1996 | 0 | 0 | 0 | 12 | 10 | 10 | 6 | 5 | 12 | 8 | 63 |
| 1997 | 0 | 0 | 0 | 24 | 28 | 4 | 6 | 2 | 5 | 5 | 74 |
| 1998 | 0 | 0 | 0 | 16 | 11 | 5 | 2 | 3 | 1 | 11 | 49 |
| 1999 | 0 | 1 | 6 | 15 | 12 | 5 | 1 | 7 | 1 | 3 | 51 |
| 2000 | 0 | 2 | 7 | 15 | 15 | 6 | 2 | 7 | 3 | 4 | 61 |
| 2001 | 0 | 0 | 1 | 12 | 15 | 2 | 1 | 2 | 2 | 7 | 42 |
| Total | 0 | 4 | 14 | 115 | 105 | 41 | 20 | 36 | 25 | 39 | 399 |

VL = very low, L = low, M = medium, H = high, VH = very high.

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1 | 10 | 13 | 9 | 3 | 1 | 6 | |
| Highest: | 330 | 517 | 394 | 184 | 863 | 846 | 640 | |
| Mean: | 64 | 120 | 72 | 133 | 81 | 78 | 107 | |
| Median: | 48 | 80 | 48 | 55 | 49 | 50 | 48 | |

Percent of home and garden samples within each Morgan extractable phosphorus range:

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 2 | 0 | 36 | 24 | 15 | 3 | 17 | 2 | 2 | 100 |
| 1996 | 0 | 0 | 0 | 19 | 16 | 16 | 10 | 8 | 19 | 13 | 100 |
| 1997 | 0 | 0 | 0 | 32 | 38 | 5 | 8 | 3 | 7 | 7 | 100 |
| 1998 | 0 | 0 | 0 | 33 | 22 | 10 | 4 | 6 | 2 | 22 | 100 |
| 1999 | 0 | 2 | 12 | 29 | 24 | 10 | 2 | 14 | 2 | 6 | 100 |
| 2000 | 0 | 3 | 11 | 25 | 25 | 10 | 3 | 11 | 5 | 7 | 100 |
| 2001 | 0 | 0 | 2 | 29 | 36 | 5 | 2 | 5 | 5 | 17 | 100 |
| Total | 0 | 1 | 4 | 29 | 26 | 10 | 5 | 9 | 6 | 10 | 100 |

VL = very low, L = low, M = medium, H = high, VH = very high.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

7.4 Richmond

Number of home and garden samples within each range Morgan extractable P range (lbs/acre Morgan P):

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 1996 | 0 | 6 | 15 | 8 | 2 | 0 | 0 | 3 | 0 | 1 | 35 |
| 1997 | 0 | 0 | 0 | 0 | 2 | 1 | 0 | 1 | 0 | 3 | 7 |
| 1998 | 0 | 0 | 0 | 4 | 0 | 0 | 0 | 0 | 2 | 0 | 6 |
| 1999 | 0 | 1 | 2 | 4 | 4 | 2 | 2 | 0 | 0 | 4 | 19 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 0 | 4 | 7 | 0 | 0 | 0 | 2 | 0 | 17 | 30 |
| Total | 0 | 7 | 21 | 25 | 8 | 3 | 2 | 6 | 2 | 26 | 100 |

VL = very low, L = low, M = medium, H = high, VH = very high.

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 29 | 1 | 41 | 13 | 2 | 441 | 4 | |
| Highest: | 31 | 336 | 404 | 167 | 1037 | 441 | 564 | |
| Mean: | 30 | 29 | 180 | 71 | 146 | 441 | 215 | |
| Median: | 30 | 7 | 127 | 37 | 52 | 441 | 227 | |

Percent of home and garden samples within each Morgan extractable phosphorus range:

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | . | . | . | . | . | . | . | . | . | . | 100 |
| 1996 | . | . | . | . | . | . | . | . | . | . | 100 |
| 1997 | . | . | . | . | . | . | . | . | . | . | 100 |
| 1998 | . | . | . | . | . | . | . | . | . | . | 100 |
| 1999 | . | . | . | . | . | . | . | . | . | . | 100 |
| 2000 | . | . | . | . | . | . | . | . | . | . | 100 |
| 2001 | . | . | . | . | . | . | . | . | . | . | 100 |
| Total | 0 | 7 | 21 | 25 | 8 | 3 | 2 | 6 | 2 | 26 | 100 |

VL = very low, L = low, M = medium, H = high, VH = very high.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

7.5 Manhattan

Number of home and garden samples within each range Morgan extractable P range (lbs/acre Morgan P):

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 3 | 2 | 17 | 6 | 2 | 5 | 8 | 4 | 9 | 56 |
| 1996 | 0 | 0 | 0 | 1 | 9 | 7 | 6 | 6 | 2 | 7 | 38 |
| 1997 | 0 | 0 | 1 | 8 | 5 | 3 | 3 | 6 | 0 | 3 | 29 |
| 1998 | 0 | 0 | 3 | 17 | 19 | 6 | 4 | 14 | 5 | 23 | 91 |
| 1999 | 0 | 4 | 5 | 19 | 14 | 13 | 9 | 5 | 4 | 13 | 86 |
| 2000 | 0 | 14 | 26 | 70 | 32 | 14 | 10 | 10 | 6 | 11 | 193 |
| 2001 | 0 | 1 | 4 | 19 | 14 | 10 | 8 | 12 | 7 | 6 | 81 |
| Total | 0 | 22 | 41 | 151 | 99 | 55 | 45 | 61 | 28 | 72 | 574 |

VL = very low, L = low, M = medium, H = high, VH = very high.

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 2 | 24 | 5 | 5 | 1 | 2 | 2 | |
| Highest: | 447 | 599 | 460 | 2427 | 1244 | 1580 | 575 | |
| Mean: | 101 | 131 | 98 | 206 | 133 | 61 | 99 | |
| Median: | 62 | 87 | 62 | 87 | 62 | 29 | 65 | |

Percent of home and garden samples within each Morgan extractable phosphorus range:

| | <1 | 1-3 | 4-8 | 9-39 | 40-60 | 61-80 | 81-100 | 101-150 | 151-200 | >200 | Total |
|-------|----|-----|-----|------|-------|-------|--------|---------|---------|------|-------|
| | VL | L | M | H | VH | VH | VH | VH | VH | VH | |
| 1995 | 0 | 5 | 4 | 30 | 11 | 4 | 9 | 14 | 7 | 16 | 100 |
| 1996 | 0 | 0 | 0 | 3 | 24 | 18 | 16 | 16 | 5 | 18 | 100 |
| 1997 | 0 | 0 | 3 | 28 | 17 | 10 | 10 | 21 | 0 | 10 | 100 |
| 1998 | 0 | 0 | 3 | 19 | 21 | 7 | 4 | 15 | 5 | 25 | 100 |
| 1999 | 0 | 5 | 6 | 22 | 16 | 15 | 10 | 6 | 5 | 15 | 100 |
| 2000 | 0 | 7 | 13 | 36 | 17 | 7 | 5 | 5 | 3 | 6 | 100 |
| 2001 | 0 | 1 | 5 | 23 | 17 | 12 | 10 | 15 | 9 | 7 | 100 |
| Total | 0 | 4 | 7 | 26 | 17 | 10 | 8 | 11 | 5 | 13 | 100 |

VL = very low, L = low, M = medium, H = high, VH = very high.

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

8. Potassium

8.1 Bronx

Number of home and garden samples within each K range (lbs K/acre Morgan extraction):

| Soil Management Group 1 | | | | | | |
|-------------------------|----------|-------|--------|---------|-----------|-------|
| | <35 | 35-64 | 65-94 | 95-149 | >149 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |
| Soil Management Group 2 | | | | | | |
| | <40 | 40-69 | 70-99 | 100-164 | >164 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 1 | 1 | 2 |
| 1997 | 0 | 0 | 1 | 1 | 6 | 8 |
| 1998 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 1 | 1 |
| Total (#) | 0 | 0 | 1 | 2 | 9 | 12 |
| Total (%) | 0 | 0 | 8 | 17 | 75 | 100 |
| Soil Management Group 3 | | | | | | |
| | <45 | 45-79 | 80-119 | 120-199 | >199 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1996 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1997 | 0 | 0 | 1 | 5 | 9 | 15 |
| 1998 | 0 | 0 | 0 | 4 | 12 | 16 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 1 | 0 | 1 | 3 | 0 | 5 |
| 2001 | 0 | 0 | 0 | 0 | 1 | 1 |
| Total (#) | 1 | 0 | 2 | 12 | 25 | 40 |
| Total (%) | 3 | 0 | 5 | 30 | 63 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

| Soil Management Group 4 | | | | | | |
|-------------------------|----------|-------|---------|---------|-----------|-------|
| | <55 | 55-99 | 100-149 | 150-239 | >239 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 3 | 12 | 16 | 12 | 43 |
| 1996 | 0 | 0 | 0 | 0 | 5 | 5 |
| 1997 | 0 | 0 | 3 | 3 | 1 | 7 |
| 1998 | 0 | 0 | 4 | 0 | 2 | 6 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 2 | 5 | 7 |
| 2001 | 0 | 0 | 0 | 1 | 0 | 1 |
| Total (#) | 0 | 3 | 19 | 22 | 25 | 69 |
| Total (%) | 0 | 4 | 28 | 32 | 36 | 100 |

| Soil Management Group 5 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 1 | 0 | 0 | 2 | 3 |
| 1997 | 0 | 0 | 0 | 2 | 5 | 7 |
| 1998 | 0 | 0 | 0 | 1 | 3 | 4 |
| 1999 | 0 | 0 | 0 | 2 | 0 | 2 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 1 | 2 | 1 | 4 |
| Total (#) | 0 | 1 | 1 | 7 | 11 | 20 |
| Total (%) | 0 | 5 | 5 | 35 | 55 | 100 |

| Soil Management Group 6 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Number of home and garden samples within each potassium classification:

| Summary (#) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 3 | 12 | 16 | 13 | 44 |
| 1996 | 0 | 1 | 0 | 1 | 10 | 12 |
| 1997 | 0 | 0 | 5 | 11 | 21 | 37 |
| 1998 | 0 | 0 | 4 | 5 | 18 | 27 |
| 1999 | 0 | 0 | 0 | 2 | 0 | 2 |
| 2000 | 1 | 0 | 1 | 5 | 5 | 12 |
| 2001 | 0 | 0 | 1 | 3 | 3 | 7 |
| Total # | 1 | 4 | 23 | 43 | 70 | 141 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 87 | 114 | 87 | 110 | 202 | 34 | 164 | |
| Highest: | 397 | 951 | 545 | 933 | 223 | 478 | 714 | |
| Mean: | 196 | 418 | 238 | 349 | 213 | 221 | 334 | |
| Median: | 177 | 330 | 218 | 310 | 213 | 209 | 222 | |

Percent of samples submitted for home and garden within each potassium classification.

| Summary (%) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 7 | 27 | 36 | 30 | 100 |
| 1996 | 0 | 8 | 0 | 8 | 83 | 100 |
| 1997 | 0 | 0 | 14 | 30 | 57 | 100 |
| 1998 | 0 | 0 | 15 | 19 | 67 | 100 |
| 1999 | 0 | 0 | 0 | 100 | 0 | 100 |
| 2000 | 8 | 0 | 8 | 42 | 42 | 100 |
| 2001 | 0 | 0 | 14 | 43 | 43 | 100 |
| Grand Total | 1 | 3 | 16 | 30 | 50 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

8.2 Queens

Number of home and garden samples within each K range (lbs K/acre Morgan extraction):

| Soil Management Group 1 | | | | | | |
|-------------------------|----------|-------|--------|---------|-----------|-------|
| | <35 | 35-64 | 65-94 | 95-149 | >149 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |
| Soil Management Group 2 | | | | | | |
| | <40 | 40-69 | 70-99 | 100-164 | >164 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 1 | 4 | 5 |
| 1996 | 0 | 0 | 0 | 1 | 2 | 3 |
| 1997 | 0 | 0 | 0 | 1 | 4 | 5 |
| 1998 | 0 | 2 | 0 | 1 | 1 | 4 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 1 | 0 | 3 | 3 | 7 |
| 2001 | 0 | 0 | 0 | 0 | 2 | 2 |
| Total (#) | 0 | 3 | 0 | 7 | 16 | 26 |
| Total (%) | 0 | 12 | 0 | 27 | 62 | 100 |
| Soil Management Group 3 | | | | | | |
| | <45 | 45-79 | 80-119 | 120-199 | >199 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 1 | 0 | 0 | 4 | 5 |
| 1996 | 0 | 0 | 2 | 4 | 14 | 20 |
| 1997 | 2 | 0 | 2 | 4 | 1 | 9 |
| 1998 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1999 | 11 | 5 | 3 | 8 | 17 | 44 |
| 2000 | 13 | 7 | 6 | 7 | 18 | 51 |
| 2001 | 0 | 0 | 0 | 2 | 7 | 9 |
| Total (#) | 26 | 13 | 13 | 25 | 63 | 140 |
| Total (%) | 19 | 9 | 9 | 18 | 45 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

| Soil Management Group 4 | | | | | | |
|-------------------------|----------|-------|---------|---------|-----------|-------|
| | <55 | 55-99 | 100-149 | 150-239 | >239 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 1 | 4 | 5 |
| 1996 | 3 | 4 | 3 | 3 | 7 | 20 |
| 1997 | 1 | 4 | 1 | 2 | 4 | 12 |
| 1998 | 0 | 0 | 3 | 7 | 3 | 13 |
| 1999 | 0 | 0 | 2 | 3 | 0 | 5 |
| 2000 | 0 | 0 | 0 | 0 | 5 | 5 |
| 2001 | 0 | 0 | 1 | 2 | 5 | 8 |
| Total (#) | 4 | 8 | 10 | 18 | 28 | 68 |
| Total (%) | 6 | 12 | 15 | 26 | 41 | 100 |

| Soil Management Group 5 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 7 | 6 | 1 | 2 | 16 |
| 1996 | 1 | 0 | 0 | 1 | 0 | 2 |
| 1997 | 15 | 4 | 5 | 3 | 3 | 30 |
| 1998 | 4 | 5 | 1 | 1 | 1 | 12 |
| 1999 | 1 | 1 | 0 | 2 | 2 | 6 |
| 2000 | 15 | 8 | 5 | 9 | 6 | 43 |
| 2001 | 0 | 1 | 0 | 3 | 4 | 8 |
| Total (#) | 36 | 26 | 17 | 20 | 18 | 117 |
| Total (%) | 31 | 22 | 15 | 17 | 15 | 100 |

| Soil Management Group 6 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Number of home and garden samples within each potassium classification:

| Summary (#) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 8 | 6 | 3 | 14 | 31 |
| 1996 | 4 | 4 | 5 | 9 | 23 | 45 |
| 1997 | 18 | 8 | 8 | 10 | 12 | 56 |
| 1998 | 4 | 7 | 4 | 9 | 7 | 31 |
| 1999 | 12 | 6 | 5 | 13 | 19 | 55 |
| 2000 | 28 | 16 | 11 | 19 | 32 | 106 |
| 2001 | 0 | 1 | 1 | 7 | 18 | 27 |
| Total # | 66 | 50 | 40 | 70 | 125 | 351 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 70 | 39 | 14 | 42 | 26 | 12 | 70 | |
| Highest: | 1002 | 1304 | 1009 | 654 | 1316 | 605 | 1866 | |
| Mean: | 228 | 246 | 162 | 173 | 194 | 156 | 393 | |
| Median: | 180 | 223 | 108 | 133 | 159 | 126 | 278 | |

Percent of samples submitted for home and garden within each potassium classification.

| Summary (%) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 26 | 19 | 10 | 45 | 100 |
| 1996 | 9 | 9 | 11 | 20 | 51 | 100 |
| 1997 | 32 | 14 | 14 | 18 | 21 | 100 |
| 1998 | 13 | 23 | 16 | 29 | 23 | 100 |
| 1999 | 22 | 11 | 9 | 24 | 35 | 100 |
| 2000 | 26 | 15 | 10 | 18 | 30 | 100 |
| 2001 | 0 | 4 | 4 | 26 | 67 | 100 |
| Grand Total | 19 | 14 | 11 | 20 | 36 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

8.3 Kings

Number of home and garden samples within each K range (lbs K/acre Morgan extraction):

| Soil Management Group 1 | | | | | | |
|-------------------------|----------|-------|--------|---------|-----------|-------|
| | <35 | 35-64 | 65-94 | 95-149 | >149 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |
| Soil Management Group 2 | | | | | | |
| | <40 | 40-69 | 70-99 | 100-164 | >164 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1996 | 0 | 0 | 1 | 0 | 17 | 18 |
| 1997 | 0 | 0 | 0 | 7 | 12 | 19 |
| 1998 | 0 | 0 | 0 | 3 | 4 | 7 |
| 1999 | 0 | 0 | 0 | 0 | 5 | 5 |
| 2000 | 0 | 1 | 0 | 0 | 2 | 3 |
| 2001 | 0 | 0 | 1 | 0 | 1 | 2 |
| Total (#) | 0 | 1 | 2 | 10 | 43 | 56 |
| Total (%) | 0 | 2 | 4 | 18 | 77 | 100 |
| Soil Management Group 3 | | | | | | |
| | <45 | 45-79 | 80-119 | 120-199 | >199 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 1 | 2 | 4 | 18 | 25 |
| 1996 | 0 | 0 | 2 | 3 | 10 | 15 |
| 1997 | 0 | 0 | 1 | 4 | 11 | 16 |
| 1998 | 0 | 1 | 3 | 3 | 19 | 26 |
| 1999 | 0 | 1 | 2 | 8 | 14 | 25 |
| 2000 | 0 | 0 | 4 | 5 | 22 | 31 |
| 2001 | 1 | 0 | 2 | 6 | 2 | 11 |
| Total (#) | 1 | 3 | 16 | 33 | 96 | 149 |
| Total (%) | 1 | 2 | 11 | 22 | 64 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

| Soil Management Group 4 | | | | | | |
|-------------------------|----------|-------|---------|---------|-----------|-------|
| | <55 | 55-99 | 100-149 | 150-239 | >239 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 8 | 6 | 12 | 26 |
| 1996 | 0 | 0 | 1 | 5 | 12 | 18 |
| 1997 | 0 | 1 | 7 | 11 | 13 | 32 |
| 1998 | 0 | 0 | 1 | 3 | 5 | 9 |
| 1999 | 0 | 2 | 1 | 2 | 6 | 11 |
| 2000 | 1 | 1 | 2 | 3 | 10 | 17 |
| 2001 | 0 | 4 | 9 | 5 | 8 | 26 |
| Total (#) | 1 | 8 | 29 | 35 | 66 | 139 |
| Total (%) | 1 | 6 | 21 | 25 | 47 | 100 |

| Soil Management Group 5 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 2 | 1 | 0 | 3 | 0 | 6 |
| 1996 | 0 | 0 | 4 | 1 | 7 | 12 |
| 1997 | 0 | 2 | 4 | 1 | 0 | 7 |
| 1998 | 0 | 2 | 0 | 3 | 2 | 7 |
| 1999 | 3 | 1 | 4 | 1 | 1 | 10 |
| 2000 | 2 | 2 | 1 | 1 | 4 | 10 |
| 2001 | 0 | 0 | 0 | 3 | 0 | 3 |
| Total (#) | 7 | 8 | 13 | 13 | 14 | 55 |
| Total (%) | 15 | 15 | 24 | 24 | 25 | 100 |

| Soil Management Group 6 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Number of home and garden samples within each potassium classification:

| Summary (#) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 2 | 2 | 10 | 13 | 32 | 59 |
| 1996 | 0 | 0 | 8 | 9 | 46 | 63 |
| 1997 | 0 | 3 | 12 | 23 | 36 | 74 |
| 1998 | 0 | 3 | 4 | 12 | 30 | 49 |
| 1999 | 3 | 4 | 7 | 11 | 26 | 51 |
| 2000 | 3 | 4 | 7 | 9 | 38 | 61 |
| 2001 | 1 | 4 | 12 | 14 | 11 | 42 |
| Total # | 9 | 20 | 60 | 91 | 219 | 399 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 51 | 89 | 66 | 75 | 33 | 24 | 27 | |
| Highest: | 817 | 1316 | 903 | 4430 | 8099 | 6931 | 784 | |
| Mean: | 255 | 337 | 237 | 629 | 375 | 465 | 219 | |
| Median: | 226 | 261 | 207 | 243 | 208 | 233 | 172 | |

Percent of samples submitted for home and garden within each potassium classification.

| Summary (%) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 3 | 3 | 17 | 22 | 54 | 100 |
| 1996 | 0 | 0 | 13 | 14 | 73 | 100 |
| 1997 | 0 | 4 | 16 | 31 | 49 | 100 |
| 1998 | 0 | 6 | 8 | 24 | 61 | 100 |
| 1999 | 6 | 8 | 14 | 22 | 51 | 100 |
| 2000 | 5 | 7 | 11 | 15 | 62 | 100 |
| 2001 | 2 | 10 | 29 | 33 | 26 | 100 |
| Grand Total | 2 | 5 | 15 | 23 | 55 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

8.4 Richmond

Number of home and garden samples within each K range (lbs K/acre Morgan extraction):

| Soil Management Group 1 | | | | | | |
|-------------------------|----------|-------|--------|---------|-----------|-------|
| | <35 | 35-64 | 65-94 | 95-149 | >149 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |
| Soil Management Group 2 | | | | | | |
| | <40 | 40-69 | 70-99 | 100-164 | >164 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 1 | 0 | 1 |
| 1996 | 0 | 0 | 0 | 1 | 4 | 5 |
| 1997 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1998 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1999 | 0 | 0 | 0 | 0 | 5 | 5 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 1 | 0 | 1 | 2 | 4 |
| Total (#) | 0 | 1 | 0 | 3 | 13 | 17 |
| Total (%) | 0 | 6 | 0 | 18 | 76 | 100 |
| Soil Management Group 3 | | | | | | |
| | <45 | 45-79 | 80-119 | 120-199 | >199 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1996 | 0 | 2 | 9 | 11 | 5 | 27 |
| 1997 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 1 | 5 | 6 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 2 | 4 | 17 | 23 |
| Total (#) | 0 | 2 | 11 | 16 | 30 | 59 |
| Total (%) | 0 | 3 | 19 | 27 | 51 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

| Soil Management Group 4 | | | | | | |
|-------------------------|----------|-------|---------|---------|-----------|-------|
| | <55 | 55-99 | 100-149 | 150-239 | >239 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1997 | 0 | 0 | 0 | 1 | 1 | 2 |
| 1998 | 0 | 2 | 0 | 2 | 1 | 5 |
| 1999 | 0 | 1 | 2 | 0 | 3 | 6 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 3 | 3 |
| Total (#) | 0 | 3 | 2 | 3 | 9 | 17 |
| Total (%) | 0 | 18 | 12 | 18 | 53 | 100 |

| Soil Management Group 5 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 2 | 0 | 2 |
| 1997 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 1 | 0 | 0 | 0 | 1 | 2 |
| 2000 | 0 | 0 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 1 | 0 | 0 | 2 | 4 | 7 |
| Total (%) | 14 | 0 | 0 | 29 | 57 | 100 |

| Soil Management Group 6 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Number of home and garden samples within each potassium classification:

| Summary (#) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 0 | 0 | 1 | 1 | 2 |
| 1996 | 0 | 2 | 9 | 14 | 10 | 35 |
| 1997 | 0 | 0 | 0 | 1 | 6 | 7 |
| 1998 | 0 | 2 | 0 | 2 | 2 | 6 |
| 1999 | 1 | 1 | 2 | 1 | 14 | 19 |
| 2000 | 0 | 0 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 1 | 2 | 5 | 22 | 30 |
| Total # | 1 | 6 | 13 | 24 | 56 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 160 | 70 | 223 | 76 | 2 | 1050 | 45 | |
| Highest: | 279 | 803 | 2133 | 967 | 7330 | 1050 | 1910 | |
| Mean: | 219 | 179 | 887 | 309 | 963 | 1050 | 836 | |
| Median: | 219 | 153 | 761 | 177 | 299 | 1050 | 515 | |

Percent of samples submitted for home and garden within each potassium classification.

| Summary (%) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 0 | 0 | 0 | 50 | 50 | 100 |
| 1996 | 0 | 6 | 26 | 40 | 29 | 100 |
| 1997 | 0 | 0 | 0 | 14 | 86 | 100 |
| 1998 | 0 | 33 | 0 | 33 | 33 | 100 |
| 1999 | 5 | 5 | 11 | 5 | 74 | 100 |
| 2000 | 0 | 0 | 0 | 0 | 100 | 100 |
| 2001 | 0 | 1 | 2 | 5 | 22 | 100 |
| Grand Total | 1 | 6 | 13 | 24 | 56 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

8.5 Manhattan

Number of home and garden samples within each K range (lbs K/acre Morgan extraction):

| Soil Management Group 1 | | | | | | |
|-------------------------|----------|-------|--------|---------|-----------|-------|
| | <35 | 35-64 | 65-94 | 95-149 | >149 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |
| Soil Management Group 2 | | | | | | |
| | <40 | 40-69 | 70-99 | 100-164 | >164 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 1 | 3 | 4 |
| 1996 | 0 | 0 | 0 | 1 | 4 | 5 |
| 1997 | 0 | 0 | 0 | 0 | 1 | 1 |
| 1998 | 0 | 0 | 0 | 5 | 10 | 15 |
| 1999 | 0 | 0 | 0 | 5 | 13 | 18 |
| 2000 | 0 | 0 | 0 | 1 | 11 | 12 |
| 2001 | 0 | 0 | 0 | 2 | 7 | 9 |
| Total (#) | 0 | 0 | 0 | 15 | 49 | 64 |
| Total (%) | 0 | 0 | 0 | 23 | 77 | 100 |
| Soil Management Group 3 | | | | | | |
| | <45 | 45-79 | 80-119 | 120-199 | >199 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 3 | 3 | 0 | 3 | 22 | 31 |
| 1996 | 0 | 1 | 1 | 2 | 9 | 13 |
| 1997 | 0 | 1 | 1 | 3 | 5 | 10 |
| 1998 | 0 | 0 | 0 | 3 | 16 | 19 |
| 1999 | 0 | 1 | 4 | 6 | 19 | 30 |
| 2000 | 0 | 1 | 2 | 17 | 30 | 50 |
| 2001 | 0 | 0 | 0 | 2 | 21 | 23 |
| Total (#) | 3 | 7 | 8 | 36 | 122 | 176 |
| Total (%) | 2 | 4 | 5 | 20 | 69 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

| Soil Management Group 4 | | | | | | |
|-------------------------|----------|-------|---------|---------|-----------|-------|
| | <55 | 55-99 | 100-149 | 150-239 | >239 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 1 | 4 | 10 | 15 |
| 1996 | 0 | 0 | 0 | 5 | 7 | 12 |
| 1997 | 0 | 0 | 1 | 3 | 8 | 12 |
| 1998 | 0 | 1 | 5 | 2 | 30 | 38 |
| 1999 | 0 | 0 | 0 | 9 | 12 | 21 |
| 2000 | 0 | 1 | 4 | 15 | 35 | 55 |
| 2001 | 0 | 1 | 1 | 9 | 23 | 34 |
| Total (#) | 0 | 3 | 12 | 47 | 125 | 187 |
| Total (%) | 0 | 2 | 6 | 25 | 67 | 100 |

| Soil Management Group 5 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 6 | 6 |
| 1996 | 0 | 0 | 3 | 2 | 3 | 8 |
| 1997 | 0 | 1 | 0 | 2 | 3 | 6 |
| 1998 | 0 | 3 | 2 | 6 | 8 | 19 |
| 1999 | 5 | 1 | 4 | 4 | 3 | 17 |
| 2000 | 51 | 3 | 2 | 7 | 13 | 76 |
| 2001 | 0 | 1 | 4 | 5 | 5 | 15 |
| Total (#) | 56 | 9 | 15 | 26 | 41 | 147 |
| Total (%) | 38 | 6 | 10 | 18 | 28 | 100 |

| Soil Management Group 6 | | | | | | |
|-------------------------|----------|--------|---------|---------|-----------|-------|
| | <60 | 60-114 | 115-164 | 165-269 | >269 | Total |
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1996 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1997 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1998 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1999 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2000 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2001 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (#) | 0 | 0 | 0 | 0 | 0 | 0 |
| Total (%) | - | - | - | - | - | - |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Number of home and garden samples within each potassium classification:

| Summary (#) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 3 | 3 | 1 | 8 | 41 | 56 |
| 1996 | 0 | 1 | 4 | 10 | 23 | 38 |
| 1997 | 0 | 2 | 2 | 8 | 17 | 29 |
| 1998 | 0 | 4 | 7 | 16 | 64 | 91 |
| 1999 | 5 | 2 | 8 | 24 | 47 | 86 |
| 2000 | 51 | 5 | 8 | 40 | 89 | 193 |
| 2001 | 0 | 2 | 5 | 18 | 56 | 81 |
| Total # | 59 | 19 | 35 | 124 | 337 | 574 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|-------|------|------|------|--|
| Lowest: | 27 | 62 | 49 | 77 | 13 | 1 | 70 | |
| Highest: | 2299 | 3016 | 622 | 17476 | 1851 | 4693 | 873 | |
| Mean: | 360 | 481 | 286 | 644 | 297 | 306 | 310 | |
| Median: | 283 | 306 | 268 | 287 | 215 | 210 | 297 | |

Percent of samples submitted for home and garden within each potassium classification.

| Summary (%) | Very Low | Low | Medium | High | Very High | Total |
|-------------|----------|-----|--------|------|-----------|-------|
| 1995 | 5 | 5 | 2 | 14 | 73 | 100 |
| 1996 | 0 | 3 | 11 | 26 | 61 | 100 |
| 1997 | 0 | 7 | 7 | 28 | 59 | 100 |
| 1998 | 0 | 4 | 8 | 18 | 70 | 100 |
| 1999 | 6 | 2 | 9 | 28 | 55 | 100 |
| 2000 | 28 | 3 | 4 | 22 | 43 | 100 |
| 2001 | 0 | 2 | 6 | 22 | 69 | 100 |
| Grand Total | 10 | 3 | 6 | 22 | 59 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

9. Magnesium

9.1 Bronx

Number of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 2 | 6 | 5 | 8 | 23 | 44 |
| 1996 | 0 | 0 | 0 | 0 | 12 | 12 |
| 1997 | 0 | 0 | 0 | 4 | 33 | 37 |
| 1998 | 0 | 0 | 1 | 2 | 24 | 27 |
| 1999 | 0 | 0 | 0 | 0 | 2 | 2 |
| 2000 | 0 | 0 | 1 | 1 | 10 | 12 |
| 2001 | 0 | 0 | 0 | 0 | 7 | 7 |
| Total | 2 | 6 | 7 | 15 | 111 | 141 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 19 | 247 | 136 | 92 | 416 | 100 | 217 | |
| Highest: | 792 | 1026 | 919 | 4432 | 426 | 854 | 648 | |
| Mean: | 262 | 477 | 404 | 1036 | 421 | 372 | 418 | |
| Median: | 203 | 427 | 376 | 486 | 421 | 312 | 442 | |

Percent of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 5 | 14 | 11 | 18 | 52 | 100 |
| 1996 | 0 | 0 | 0 | 0 | 100 | 100 |
| 1997 | 0 | 0 | 0 | 11 | 89 | 100 |
| 1998 | 0 | 0 | 4 | 7 | 89 | 100 |
| 1999 | 0 | 0 | 0 | 0 | 100 | 100 |
| 2000 | 0 | 0 | 8 | 8 | 83 | 100 |
| 2001 | 0 | 0 | 0 | 0 | 100 | 100 |
| Total | 1 | 4 | 5 | 11 | 79 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

9.2 Queens

Number of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 1 | 0 | 6 | 24 | 31 |
| 1996 | 0 | 1 | 3 | 13 | 28 | 45 |
| 1997 | 6 | 9 | 3 | 5 | 33 | 56 |
| 1998 | 0 | 1 | 3 | 5 | 22 | 31 |
| 1999 | 3 | 12 | 3 | 6 | 31 | 55 |
| 2000 | 6 | 23 | 4 | 24 | 49 | 106 |
| 2001 | 0 | 0 | 1 | 1 | 25 | 27 |
| Total | 15 | 47 | 17 | 60 | 212 | 351 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 58 | 65 | 11 | 52 | 13 | 15 | 91 | |
| Highest: | 2359 | 1531 | 996 | 1431 | 967 | 938 | 1149 | |
| Mean: | 412 | 338 | 242 | 331 | 267 | 232 | 545 | |
| Median: | 301 | 259 | 238 | 274 | 229 | 177 | 415 | |

Percent of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 3 | 0 | 19 | 77 | 100 |
| 1996 | 0 | 2 | 7 | 29 | 62 | 100 |
| 1997 | 11 | 16 | 5 | 9 | 59 | 100 |
| 1998 | 0 | 3 | 10 | 16 | 71 | 100 |
| 1999 | 5 | 22 | 5 | 11 | 56 | 100 |
| 2000 | 6 | 22 | 4 | 23 | 46 | 100 |
| 2001 | 0 | 0 | 4 | 4 | 93 | 100 |
| Total | 4 | 13 | 5 | 17 | 60 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

9.3 Kings

Number of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 1 | 1 | 16 | 41 | 59 |
| 1996 | 0 | 0 | 0 | 3 | 60 | 63 |
| 1997 | 0 | 1 | 3 | 22 | 48 | 74 |
| 1998 | 0 | 1 | 0 | 9 | 39 | 49 |
| 1999 | 0 | 0 | 3 | 8 | 40 | 51 |
| 2000 | 0 | 5 | 3 | 6 | 47 | 61 |
| 2001 | 0 | 2 | 0 | 8 | 32 | 42 |
| Total | 0 | 10 | 10 | 72 | 307 | 399 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 50 | 101 | 61 | 46 | 77 | 51 | 42 | |
| Highest: | 740 | 2167 | 2291 | 5386 | 2513 | 3584 | 1684 | |
| Mean: | 307 | 500 | 358 | 627 | 370 | 446 | 495 | |
| Median: | 289 | 343 | 264 | 361 | 281 | 316 | 304 | |

Percent of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 2 | 2 | 27 | 69 | 100 |
| 1996 | 0 | 0 | 0 | 5 | 95 | 100 |
| 1997 | 0 | 1 | 4 | 30 | 65 | 100 |
| 1998 | 0 | 2 | 0 | 18 | 80 | 100 |
| 1999 | 0 | 0 | 6 | 16 | 78 | 100 |
| 2000 | 0 | 8 | 5 | 10 | 77 | 100 |
| 2001 | 0 | 5 | 0 | 19 | 76 | 100 |
| Total | 0 | 3 | 3 | 18 | 77 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

9.4 Richmond

Number of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 0 | 0 | 0 | 2 | 2 |
| 1996 | 0 | 0 | 0 | 1 | 34 | 35 |
| 1997 | 0 | 0 | 0 | 0 | 7 | 7 |
| 1998 | 0 | 0 | 0 | 2 | 4 | 6 |
| 1999 | 0 | 1 | 2 | 2 | 14 | 19 |
| 2000 | 0 | 0 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 0 | 1 | 1 | 28 | 30 |
| Total | 0 | 1 | 3 | 6 | 90 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 471 | 180 | 477 | 148 | 29 | 847 | 69 | |
| Highest: | 513 | 1095 | 2144 | 1133 | 3855 | 847 | 1132 | |
| Mean: | 492 | 447 | 861 | 503 | 626 | 847 | 402 | |
| Median: | 492 | 408 | 737 | 429 | 416 | 847 | 375 | |

Percent of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | . | . | . | . | . | 100 |
| 1996 | . | . | . | . | . | 100 |
| 1997 | . | . | . | . | . | 100 |
| 1998 | . | . | . | . | . | 100 |
| 1999 | . | . | . | . | . | 100 |
| 2000 | . | . | . | . | . | 100 |
| 2001 | . | . | . | . | . | 100 |
| Total | 0 | 1 | 3 | 6 | 90 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

9.5 Manhattan

Number of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 4 | 2 | 7 | 43 | 56 |
| 1996 | 0 | 2 | 1 | 8 | 27 | 38 |
| 1997 | 0 | 1 | 1 | 3 | 24 | 29 |
| 1998 | 0 | 1 | 3 | 12 | 75 | 91 |
| 1999 | 1 | 0 | 3 | 25 | 57 | 86 |
| 2000 | 2 | 47 | 5 | 16 | 123 | 193 |
| 2001 | 0 | 2 | 2 | 12 | 65 | 81 |
| Total | 3 | 57 | 17 | 83 | 414 | 574 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 21 | 47 | 59 | 61 | 18 | 18 | 27 | |
| Highest: | 5813 | 3294 | 975 | 6107 | 5167 | 4693 | 1585 | |
| Mean: | 491 | 659 | 356 | 665 | 834 | 362 | 449 | |
| Median: | 341 | 377 | 318 | 408 | 345 | 301 | 374 | |

Percent of home and garden samples within each Mg range (lbs Morgan Mg/acre):

| | <20 | 20-65 | 66-100 | 101-199 | >199 | Total |
|-------|----------|-------|--------|---------|-----------|-------|
| | Very Low | Low | Medium | High | Very High | |
| 1995 | 0 | 7 | 4 | 13 | 77 | 100 |
| 1996 | 0 | 5 | 3 | 21 | 71 | 100 |
| 1997 | 0 | 3 | 3 | 10 | 83 | 100 |
| 1998 | 0 | 1 | 3 | 13 | 82 | 100 |
| 1999 | 1 | 0 | 3 | 29 | 66 | 100 |
| 2000 | 1 | 24 | 3 | 8 | 64 | 100 |
| 2001 | 0 | 2 | 2 | 15 | 80 | 100 |
| Total | 1 | 10 | 3 | 14 | 72 | 100 |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

10. Iron

10.1 Bronx

Iron (lbs Fe/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-49 | >49 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 24 | 20 | 44 |
| 1996 | 8 | 4 | 12 |
| 1997 | 33 | 4 | 37 |
| 1998 | 18 | 9 | 27 |
| 1999 | 2 | 0 | 2 |
| 2000 | 10 | 2 | 12 |
| 2001 | 7 | 0 | 7 |
| Total | 102 | 39 | 141 |

Percentages:

| 0-49 | >49 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 55 | 45 | 100 |
| 67 | 33 | 100 |
| 89 | 11 | 100 |
| 67 | 33 | 100 |
| 100 | 0 | 100 |
| 83 | 17 | 100 |
| 100 | 0 | 100 |
| 72 | 28 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 3 | 2 | 2 | 3 | 6 | 2 | 2 | |
| Highest: | 639 | 71 | 794 | 205 | 8 | 53 | 10 | |
| Mean: | 68 | 31 | 52 | 50 | 7 | 13 | 7 | |
| Median: | 45 | 24 | 9 | 25 | 7 | 6 | 8 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

10.2 Queens

Iron (lbs Fe/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-49 | >49 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 30 | 1 | 31 |
| 1996 | 43 | 2 | 45 |
| 1997 | 55 | 1 | 56 |
| 1998 | 30 | 1 | 31 |
| 1999 | 50 | 5 | 55 |
| 2000 | 104 | 2 | 106 |
| 2001 | 20 | 7 | 27 |
| Total | 332 | 19 | 351 |

Percentages:

| 0-49 | >49 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 97 | 3 | 100 |
| 96 | 4 | 100 |
| 98 | 2 | 100 |
| 97 | 3 | 100 |
| 91 | 9 | 100 |
| 98 | 2 | 100 |
| 74 | 26 | 100 |
| 95 | 5 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 2 | 1 | 2 | 1 | 3 | 1 | 1 | |
| Highest: | 91 | 94 | 335 | 61 | 269 | 69 | 401 | |
| Mean: | 12 | 16 | 17 | 10 | 28 | 12 | 46 | |
| Median: | 6 | 11 | 8 | 7 | 9 | 7 | 9 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

10.3 Kings

Iron (lbs Fe/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-49 | >49 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 56 | 3 | 59 |
| 1996 | 61 | 2 | 63 |
| 1997 | 70 | 4 | 74 |
| 1998 | 46 | 3 | 49 |
| 1999 | 48 | 3 | 51 |
| 2000 | 57 | 4 | 61 |
| 2001 | 41 | 1 | 42 |
| Total | 379 | 20 | 399 |

Percentages:

| 0-49 | >49 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 95 | 5 | 100 |
| 97 | 3 | 100 |
| 95 | 5 | 100 |
| 94 | 6 | 100 |
| 94 | 6 | 100 |
| 93 | 7 | 100 |
| 98 | 2 | 100 |
| 95 | 5 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 2 | 2 | 2 | 2 | 3 | 2 | 2 | |
| Highest: | 95 | 188 | 70 | 196 | 156 | 81 | 60 | |
| Mean: | 17 | 14 | 16 | 19 | 16 | 13 | 10 | |
| Median: | 10 | 9 | 10 | 10 | 10 | 8 | 7 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

10.4 Richmond

Iron (lbs Fe/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-49 | >49 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 1 | 1 | 2 |
| 1996 | 35 | 0 | 35 |
| 1997 | 7 | 0 | 7 |
| 1998 | 6 | 0 | 6 |
| 1999 | 16 | 3 | 19 |
| 2000 | 1 | 0 | 1 |
| 2001 | 30 | 0 | 30 |
| Total | 96 | 4 | 100 |

Percentages:

| 0-49 | >49 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| 96 | 4 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 3 | 2 | 3 | 6 | 3 | 6 | 2 | |
| Highest: | 93 | 29 | 16 | 45 | 146 | 6 | 23 | |
| Mean: | 48 | 7 | 8 | 24 | 27 | 6 | 5 | |
| Median: | 48 | 5 | 7 | 23 | 10 | 6 | 4 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

10.5 Manhattan

Iron (lbs Fe/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-49 | >49 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 56 | 0 | 56 |
| 1996 | 38 | 0 | 38 |
| 1997 | 26 | 3 | 29 |
| 1998 | 85 | 6 | 91 |
| 1999 | 83 | 3 | 86 |
| 2000 | 191 | 2 | 193 |
| 2001 | 79 | 2 | 81 |
| Total | 558 | 16 | 574 |

Percentages:

| 0-49 | >49 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 90 | 10 | 100 |
| 93 | 7 | 100 |
| 97 | 3 | 100 |
| 99 | 1 | 100 |
| 98 | 2 | 100 |
| 97 | 3 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 3 | 3 | 3 | 2 | 1 | 1 | 1 | |
| Highest: | 41 | 49 | 106 | 323 | 1832 | 74 | 111 | |
| Mean: | 11 | 13 | 20 | 21 | 33 | 13 | 11 | |
| Median: | 8 | 8 | 10 | 10 | 8 | 10 | 6 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

11. Manganese

11.1 Bronx

Manganese (lbs Mn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 44 | 0 | 44 |
| 1996 | 11 | 1 | 12 |
| 1997 | 33 | 4 | 37 |
| 1998 | 21 | 6 | 27 |
| 1999 | 2 | 0 | 2 |
| 2000 | 12 | 0 | 12 |
| 2001 | 7 | 0 | 7 |
| Total | 130 | 11 | 141 |

Percentages:

| 0-99 | >99 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 100 | 0 | 100 |
| 92 | 8 | 100 |
| 89 | 11 | 100 |
| 78 | 22 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 92 | 8 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 8 | 14 | 9 | 4 | 22 | 6 | 15 | |
| Highest: | 89 | 344 | 68 | 183 | 29 | 47 | 41 | |
| Mean: | 25 | 77 | 26 | 39 | 26 | 21 | 26 | |
| Median: | 20 | 67 | 22 | 31 | 26 | 22 | 23 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

11.2 Queens

Manganese (lbs Mn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 31 | 0 | 31 |
| 1996 | 45 | 0 | 45 |
| 1997 | 56 | 0 | 56 |
| 1998 | 31 | 0 | 31 |
| 1999 | 55 | 0 | 55 |
| 2000 | 106 | 0 | 106 |
| 2001 | 25 | 2 | 27 |
| Total | 349 | 2 | 351 |

Percentages:

| 0-99 | >99 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 100 | 0 | 100 |
| 93 | 7 | 100 |
| 99 | 1 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 2 | 3 | 2 | 6 | 2 | 1 | 8 | |
| Highest: | 89 | 54 | 94 | 64 | 95 | 78 | 188 | |
| Mean: | 19 | 19 | 17 | 16 | 19 | 14 | 37 | |
| Median: | 13 | 16 | 12 | 14 | 11 | 11 | 25 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

11.3 Kings

Manganese (lbs Mn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 58 | 1 | 59 |
| 1996 | 62 | 1 | 63 |
| 1997 | 74 | 0 | 74 |
| 1998 | 46 | 3 | 49 |
| 1999 | 49 | 2 | 51 |
| 2000 | 60 | 1 | 61 |
| 2001 | 42 | 0 | 42 |
| Total | 391 | 8 | 399 |

Percentages:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 98 | 2 | 100 |
| 1996 | 98 | 2 | 100 |
| 1997 | 100 | 0 | 100 |
| 1998 | 94 | 6 | 100 |
| 1999 | 96 | 4 | 100 |
| 2000 | 98 | 2 | 100 |
| 2001 | 100 | 0 | 100 |
| Total | 98 | 2 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 3 | 5 | 4 | 8 | 4 | 3 | 4 | |
| Highest: | 125 | 101 | 55 | 663 | 126 | 110 | 64 | |
| Mean: | 18 | 25 | 20 | 43 | 24 | 23 | 21 | |
| Median: | 13 | 23 | 17 | 20 | 18 | 19 | 16 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

11.4 Richmond

Manganese (lbs Mn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 1 | 1 | 2 |
| 1996 | 35 | 0 | 35 |
| 1997 | 7 | 0 | 7 |
| 1998 | 6 | 0 | 6 |
| 1999 | 18 | 1 | 19 |
| 2000 | 1 | 0 | 1 |
| 2001 | 16 | 14 | 30 |
| Total | 84 | 16 | 100 |

Percentages:

| 0-99 | >99 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| . | . | 100 |
| 84 | 16 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 19 | 10 | 23 | 18 | 5 | 24 | 11 | |
| Highest: | 473 | 64 | 92 | 54 | 133 | 24 | 174 | |
| Mean: | 246 | 18 | 45 | 32 | 37 | 24 | 79 | |
| Median: | 246 | 16 | 33 | 24 | 38 | 24 | 56 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

11.5 Manhattan

Manganese (lbs Mn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | 0-99 | >99 | Total |
|-------|--------|-----------|-------|
| | Normal | Excessive | |
| 1995 | 55 | 1 | 56 |
| 1996 | 37 | 1 | 38 |
| 1997 | 28 | 1 | 29 |
| 1998 | 90 | 1 | 91 |
| 1999 | 84 | 2 | 86 |
| 2000 | 192 | 1 | 193 |
| 2001 | 80 | 1 | 81 |
| Total | 566 | 8 | 574 |

Percentages:

| 0-99 | >99 | Total |
|--------|-----------|-------|
| Normal | Excessive | |
| 98 | 2 | 100 |
| 97 | 3 | 100 |
| 97 | 3 | 100 |
| 99 | 1 | 100 |
| 98 | 2 | 100 |
| 99 | 1 | 100 |
| 99 | 1 | 100 |
| 99 | 1 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|------|------|------|--|
| Lowest: | 1 | 3 | 6 | 7 | 1 | 1 | 6 | |
| Highest: | 119 | 156 | 117 | 110 | 290 | 192 | 71 | |
| Mean: | 26 | 28 | 26 | 27 | 28 | 17 | 24 | |
| Median: | 21 | 19 | 19 | 20 | 18 | 14 | 22 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

12. Zinc

12.1 Bronx

Zinc (lbs Zn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 0 | 44 | 44 |
| 1996 | 0 | 0 | 12 | 12 |
| 1997 | 0 | 0 | 37 | 37 |
| 1998 | 0 | 0 | 27 | 27 |
| 1999 | 0 | 0 | 2 | 2 |
| 2000 | 0 | 0 | 12 | 12 |
| 2001 | 0 | 0 | 7 | 7 |
| Total | 0 | 0 | 141 | 141 |

Percentages:

| <0.5 | 0.5-1.0 | >1 | Total |
|------|---------|------|-------|
| Low | Medium | High | |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|-------|-------|-------|------|-------|-------|--|
| Lowest: | 2.5 | 7.6 | 1.8 | 3.5 | 7.8 | 2.4 | 9.5 | |
| Highest: | 90.1 | 128.2 | 114.0 | 226.9 | 10.3 | 127.7 | 141.8 | |
| Mean: | 17.1 | 44.0 | 29.4 | 35.9 | 9.1 | 29.2 | 37.4 | |
| Median: | 10.4 | 29.8 | 22.3 | 18.8 | 9.1 | 9.9 | 13.7 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

12.2 Queens

Zinc (lbs Zn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 0 | 31 | 31 |
| 1996 | 0 | 0 | 45 | 45 |
| 1997 | 0 | 6 | 50 | 56 |
| 1998 | 0 | 0 | 31 | 31 |
| 1999 | 0 | 2 | 53 | 55 |
| 2000 | 0 | 2 | 104 | 106 |
| 2001 | 0 | 0 | 27 | 27 |
| Total | 0 | 10 | 341 | 351 |

Percentages:

| <0.5 | 0.5-1.0 | >1 | Total |
|------|---------|------|-------|
| Low | Medium | High | |
| 0 | 0 | 100 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 11 | 89 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 4 | 96 | 100 |
| 0 | 2 | 98 | 100 |
| 0 | 0 | 100 | 100 |
| 0 | 3 | 97 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|-------|-------|-------|-------|-------|-------|-------|--|
| Lowest: | 1.9 | 2.0 | 0.7 | 1.3 | 0.7 | 0.1 | 6.0 | |
| Highest: | 178.4 | 425.5 | 252.3 | 209.3 | 210.3 | 228.3 | 443.9 | |
| Mean: | 42.2 | 77.9 | 33.1 | 34.8 | 31.2 | 24.4 | 101.3 | |
| Median: | 30.5 | 26.4 | 14.5 | 16.9 | 12.1 | 12.9 | 33.3 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

12.3 Kings

Zinc (lbs Zn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 0 | 59 | 59 |
| 1996 | 0 | 0 | 63 | 63 |
| 1997 | 0 | 0 | 74 | 74 |
| 1998 | 0 | 0 | 49 | 49 |
| 1999 | 0 | 0 | 51 | 51 |
| 2000 | 0 | 3 | 58 | 61 |
| 2001 | 0 | 0 | 42 | 42 |
| Total | 0 | 3 | 396 | 399 |

Percentages:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 0 | 100 | 100 |
| 1996 | 0 | 0 | 100 | 100 |
| 1997 | 0 | 0 | 100 | 100 |
| 1998 | 0 | 0 | 100 | 100 |
| 1999 | 0 | 0 | 100 | 100 |
| 2000 | 0 | 5 | 95 | 100 |
| 2001 | 0 | 0 | 100 | 100 |
| Total | 0 | 1 | 99 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|-------|-------|-------|-------|-------|-------|-------|--|
| Lowest: | 4.1 | 1.3 | 3.0 | 7.0 | 2.7 | 0.5 | 3.3 | |
| Highest: | 326.6 | 504.2 | 553.0 | 440.7 | 445.4 | 292.4 | 268.5 | |
| Mean: | 39.4 | 74.0 | 58.5 | 66.8 | 62.2 | 45.3 | 51.4 | |
| Median: | 21.8 | 40.1 | 34.3 | 43.0 | 25.4 | 32.2 | 39.4 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

12.4 Richmond

Zinc (lbs Zn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 0 | 2 | 2 |
| 1996 | 0 | 1 | 34 | 35 |
| 1997 | 0 | 0 | 7 | 7 |
| 1998 | 0 | 0 | 6 | 6 |
| 1999 | 0 | 0 | 19 | 19 |
| 2000 | 0 | 0 | 1 | 1 |
| 2001 | 0 | 0 | 30 | 30 |
| Total | 0 | 1 | 99 | 100 |

Percentages:

| <0.5 | 0.5-1.0 | >1 | Total |
|------|---------|------|-------|
| Low | Medium | High | |
| . | . | . | 100 |
| . | . | . | 100 |
| . | . | . | 100 |
| . | . | . | 100 |
| . | . | . | 100 |
| . | . | . | 100 |
| . | . | . | 100 |
| 0 | 1 | 99 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|------|------|------|------|-------|------|------|--|
| Lowest: | 3.3 | 0.9 | 22.4 | 5.9 | 7.3 | 32.2 | 2.2 | |
| Highest: | 3.3 | 65.8 | 74.6 | 39.8 | 139.9 | 32.2 | 65.2 | |
| Mean: | 3.3 | 8.0 | 36.7 | 21.3 | 22.5 | 32.2 | 35.1 | |
| Median: | 3.3 | 2.9 | 31.0 | 21.6 | 15.4 | 32.2 | 43.8 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

12.5 Manhattan

Zinc (lbs Zn/acre Morgan extraction) in samples for home and garden:

Total number of samples:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 3 | 53 | 56 |
| 1996 | 0 | 0 | 38 | 38 |
| 1997 | 0 | 0 | 29 | 29 |
| 1998 | 0 | 0 | 91 | 91 |
| 1999 | 0 | 0 | 86 | 86 |
| 2000 | 0 | 5 | 188 | 193 |
| 2001 | 0 | 2 | 79 | 81 |
| Total | 0 | 10 | 564 | 574 |

Percentages:

| | <0.5 | 0.5-1.0 | >1 | Total |
|-------|------|---------|------|-------|
| | Low | Medium | High | |
| 1995 | 0 | 5 | 95 | 100 |
| 1996 | 0 | 0 | 100 | 100 |
| 1997 | 0 | 0 | 100 | 100 |
| 1998 | 0 | 0 | 100 | 100 |
| 1999 | 0 | 0 | 100 | 100 |
| 2000 | 0 | 3 | 97 | 100 |
| 2001 | 0 | 2 | 98 | 100 |
| Total | 0 | 2 | 98 | 100 |

| | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | |
|----------|-------|-------|-------|-------|-------|-------|-------|--|
| Lowest: | 0.7 | 7.0 | 5.5 | 4.7 | 1.1 | 0.7 | 0.6 | |
| Highest: | 220.3 | 157.3 | 407.4 | 551.5 | 230.7 | 320.0 | 142.7 | |
| Mean: | 38.2 | 45.3 | 52.4 | 50.9 | 40.5 | 22.6 | 26.3 | |
| Median: | 15.8 | 39.5 | 19.8 | 29.7 | 24.3 | 11.1 | 15.5 | |

Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.



Ketterings, Q.M., H. Krol, C.P. Mazza, and W.S. Reid (2004). Home and community garden soil samples survey of New York City. CSS Extension Bulletin E04-21. 71 pages.

Appendix: Cornell Crop Codes

Crop codes are used in the Cornell Nutrient Analyses Laboratory.

| Crop Code | Crop Description |
|-----------|--------------------------------------|
| ALG | Ornamentals adapted to pH 4.5 to 6.0 |
| ATF | Athletic Field |
| BLU | Blueberries |
| FLA | Flowering Annuals |
| GRA | Grapes |
| HRB | Herbs |
| IDL | Idle land |
| LAW | Lawn |
| MVG | Mixed vegetables |
| OTH | Other |
| PER | Perennials |
| PRK | Park |
| ROD | Roadside |
| ROS | Roses |
| RSP | Raspberries |
| SAG | Ornamentals adapted to pH 6.0 to 7.5 |
| SPB | Spring flowering bulbs |
| STR | Strawberries |
| SUB | Summer flowering bulbs |
| TOM | Tomatoes |
| TRF | Tree fruits |